

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 308 885
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88115462.9

(51) Int. Cl. 4: **C07D 209/42** , **C07D 213/56** ,
C07D 215/16 , **C07D 241/44** ,
C07D 403/12 , **A61K 31/395**

(22) Date of filing: 21.09.88

(30) Priority: **21.09.87 US 99866**
22.08.88 US 234525

(43) Date of publication of application:
29.03.89 Bulletin 89/13

(84) Designated Contracting States:
ES GR

(71) Applicant: **ABBOTT LABORATORIES**


Abbott Park Illinois 60064(US)

(72) Inventor: **Nadzan, Alex M.**
1690 Young Drive
Libertyville Illinois 60048(US)
Inventor: **Lin, Chun Wei**
344 S. Edgewood
Wood Dale Illinois 60191(US)
Inventor: **Kerwin, James F., Jr.**
1301 Hampton Lane
Mundelein Illinois 60060(US)

(74) Representative: **Modiano, Guido et al**
MODIANO, JOSIF, PISANTY & STAUB
Modiano & Associati Via Meravigli, 16
I-20123 Milan(IT)

(54) Derivatives of d-glutamic acid and d-aspartic acid.

(57) A compound of the formula:



wherein R_1 and R_2 are substituents; R_6 is hydrogen, loweralkyl, cycloalkyl, loweralkenyl, arylalkyl, (substituted aryl)alkyl, or (heterocyclic)alkyl; B is a bond, an alkylene group, an alkenylene group, an alkadienylene group or a heteroatom containing linking group; D is a substituted acyl group or tetrazolyl; Z is CO, C(S) or S(O)₂; Ar is a heterocyclic group; and n is 1 to 3.

Also disclosed are a composition and a method for antagonizing CCK, a composition and method for treating

EP 0 308 885 A1

or preventing gastrointestinal, central nervous, appetite regulating and pain regulating systems, a composition and method for treating or preventing hyperinsulinemia, methods of making the compounds and intermediates useful in making the compounds.

DERIVATIVES OF D-GLUTAMIC ACID AND D-ASPARTIC ACID

Technical Field

This is a continuation-in-part of U.S. patent application Serial No. 099,866, filed September 21, 1987.

The present invention relates to novel organic compounds and compositions which antagonize cholecystokinin, processes for making such compounds, synthetic intermediates employed in these processes, a composition and a method for antagonizing CCK, and a composition and a method for treating gastrointestinal disorders, cancers of the gall bladder and pancreas, hyperinsulinemia, central nervous system disorders, or potentiating pain, or regulating appetite with such compounds.

10

Background of the Invention

Cholecystokinins (CCK) are a family of amino acid polypeptide hormones. CCK₃₃, a 33 amino acid fragment of CCK was first isolated from hog intestine. (Matt and Jorpes, *Biochem. J.* 125 628 (1981)). Recently the CCK₃₃ fragment has been found in the brain, where it appears to be the precursor of two smaller fragments, an octapeptide CCK₈ and a tetrapeptide CCK₄. (Dockray, *Nature* 264 4022 (1979)).

CCK₈, the carboxyl terminal octapeptide fragment of CCK, is the most potent peptide in the CCK family and is the predominant form of CCK in the brain. (Larson and Rehfeld, *Brain Res.* 165 41 (1981)). The localization of CCK fragments in the cortex of the brain suggests that CCK may be an important neuromodulator of memory, learning and control of primary sensory and motor functions. CCK and its fragments are believed to play an important role in appetite regulation and satiety. (Della-Fera, *Science* 206 471 (1979); Gibbs et al., *Nature* 289 599(1981); and Smith, *Eating and Its Disorders*, eds., Raven Press, New York, 67 (1984)).

CCK antagonists are useful in the treatment and prevention of CCK-related disorders of the gastrointestinal (GI), central nervous (CNS) and appetite regulatory systems of animals, especially man. CCK antagonists are also useful in potentiating and prolonging opiate induced analgesia and thus have utility in the treatment of pain. (Faris et al., *Science* 226 1215 (1984)).

Previously four distinct chemical classes of CCK receptor antagonists have been reported. The first class comprises derivatives of cyclic nucleotides as represented by dibutyl cyclic GMP (N. Barlos et al., *Am. J. Physiol.*, 242, G 161 (1982) and references therein). The second class is represented by C-terminal fragments of CCK (see Jensen et al., *Biochem. Biophys. Acta*, 757, 250 (1983) and Spanarkel *J. Biol. Chem.* 258, 6746 (1983)). The third class comprises amino acid derivatives of glutamic acid and tryptophan as indicated by proglumide and benzotript (see Hans et al. *Proc. Natl. Acad. Sci. U.S.A.*, 78, 6304 (1981) and Jensen et al. *Biochem. Biophys. Acta*, 761, 269 (1983)). The fourth and most recent class is comprised of 3-substituted benzodiazepines, represented by L-364,718 (see: Evans et al. *Proc. Natl. Acad. Sci. U.S.A.*, 83 4918 (1986)).

With the exception of the benzodiazepine based class, all of these compounds are relatively weak antagonists of CCK demonstrating IC₅₀ values between 10⁻⁴ and 10⁻⁵M. The benzodiazepine CCK antagonists or their metabolites may have undesirable effects *in vivo* due to their interaction with benzodiazepine receptors.

The C-terminal pentapeptide fragment of CCK is the same as the C-terminal pentapeptide fragment of another polypeptide hormone, gastrin. Gastrin, like CCK, exists in both the GI and CNS systems. Gastrin antagonists are useful in the treatment and prevention of gastrin-related disorders of the GI system such as ulcers, Zollinger-Ellison syndrome and central G cell hyperplasia. There are no known effective receptor antagonists of the *in vivo* effects of gastrin. (Morely, *Gut Pept. Ulcer Proc.*, Hiroshima Symp. 2nd, 1, (1983)).


Summary of the Invention

It has now been found that the compounds of the invention are antagonists of cholecystokinin (CCK) and bind specifically to CCK receptors. These CCK antagonists are useful in the treatment and prevention of CCK-related disorders of the gastrointestinal, central nervous and appetite regulatory systems of animals and humans. These compounds are useful in the treatment and prevention of gastrointestinal ulcers, cancers of the gall bladder and pancreas, pancreatitis, hyperinsulinemia, Zollinger-Ellison syndrome, central G cell hyperplasia, irritable bowel syndrome, the treatment or prevention of neuroleptic disorders, tardive

dyskinesia, Parkinson's disease, psychosis, Gilles de la Tourette syndrome, disorders of appetite regulatory systems, the treatment of pain and the treatment of substance abuse.

5 Disclosure of the Invention

In accordance with the present invention, there are cholecystokinin antagonists of the formula:




Formula I

wherein

R₁ and R₂ are independently selected from

- 1) hydrogen,
- 2) C₁-C₈ loweralkyl,
- 3) cycloalkyl,
- 4) loweralkenyl,
- 5) -(CH₂)_m(C(O))_rcycloalkyl wherein m and r are not both 0,
- 6) -(CH₂)_mCN,
- 7) -(CH₂)_mOR₃ wherein R₃ is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH(loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,
- 8) adamantyl,
- 9) -(CH₂)_m(C(O))_rNR₆R₇ wherein R₆ and R₇ are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, -(CH₂)_maryl, -(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, and -(CH₂)_mheterocyclic, 10) cyclic groups wherein R₁ and R₂ taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from (CH₂)_p, (CH=CH)_r, CH₂C(O), C(O)CH₂, QCH₂, CH₂Q, C(O)Q, and QC(O) wherein Q is S, O, or NR₁₀ wherein R₁₀ is hydrogen, -(C(O))_r(C₁-C₈loweralkyl), -(C(O))_rcycloalkyl, -(C(O))_rloweralkenyl, -(C(O))_rloweralkynyl, thioalkoxy, -(C(O))_r(CH₂)_maryl, -(C(O))_r(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mcarboxyl, -(CH₂)_s(C(O))_raryl, -(CH₂)_s(C(O))_r(substituted aryl), -(CH₂)_mcarboalkoxy, -(CH₂)_mcarboxamide, -(CH₂)_mcarboaryloxy, -C(X)₃ wherein X is halogen, -(C(O))_r(CH₂)_sR₁₁ wherein R₁₁ is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, -(CH₂)_maryl, --(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mheterocyclic, nitro, halogen, -CN, -OH, -NH₂, -NHOH, -NR₆R₇ wherein R₆ and R₇ are as previously defined, -NR₄OH wherein R₄ is hydrogen, -(C(O))_rloweralkyl, -(C(O))_rloweralkenyl, -(C(O))_rcycloalkyl, -(C(O))_rloweralkynyl, -(C(O))_r(CH₂)_maryl, -(C(O))_r(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mOR₄ wherein R₄ is as

defined above, or $-(CH_2)_mSR_4$ wherein R_4 is as defined above; Y is S, O, CH_2 or NR_{10} ; and R_5 is one, two or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_8 \text{ loweralkyl})$, $-(C(O))_r \text{ cycloalkyl}$, decahydronaphthyl, $-(C(O))_r \text{ loweralkenyl}$, $-(C(O))_r \text{ loweralkynyl}$, thioalkoxy, $-(C(O))_r(CH_2)_m \text{ aryl}$, $-(C(O))_r(CH_2)_m \text{ (substituted aryl)}$ wherein substituted aryl is as defined above, $-(CH_2)_s(C(O)) \text{ aryl}$, $-(CH_2)_s(C(O)) \text{ (substituted aryl)}$ wherein substituted aryl is as defined above, $-(CH_2)_m \text{ carboxyl}$, $-(CH_2)_m \text{ carboalkoxy}$, $-(CH_2)_m \text{ carboxamide}$, $-(CH_2)_m \text{ carboaryloxy}$, $-C(X)_3$, $-(C(O))_r(CH_2)_m R_6$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-(CH_2)_mOR_4$, $-(CH_2)_mSR_4$ and $-NR_4OH$ wherein R_4 is as defined above;

11) $-(C(R_5)_2)_m \text{ aryl}$ wherein R_5 is as defined above with the proviso that not all R_5 substituents are hydrogen,

12) $-(C(R_5)_2)_m \text{ (substituted aryl)}$ wherein R_5 is defined as above with the proviso that not all R_5 substituents are hydrogen and wherein substituted aryl is as defined above, and


13) $-(C(R_5)_2)_mC(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

B is

- 1) $(CH_2)_m$,
- 2) $(CR_5 = CR_5)_q$ wherein R_5 is as defined above,
- 3) LCH_2 wherein L is O, S, or NR_5 wherein R_5 is as defined above,
- 4) CH_2L wherein L is as defined above, or
- 5) NR_5 wherein R_5 is as defined above;

D is

- 1) $-C(O)R_3$ wherein R_3 is as defined above or
- 2)



Z is

- 1) $C(O)$,
- 2) $C(S)$, or
- 3) $S(O)_2$;

Ar is a heterocyclic group, n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or pharmaceutically acceptable salts thereof.

The compounds of the invention wherein $n=1-3$, D is $-C(O)R_3$ wherein R_3 is $-OH$ or $-NR_6R_7$ wherein R_6 and R_7 are independently selected from hydrogen, loweralkyl, cycloalkyl, C_2-4 loweralkenyl, benzyl, phenethyl and naphthylmethyl; R_1 and R_2 are independently selected from hydrogen, loweralkyl, cycloalkyl, C_2-4 loweralkenyl, benzyl, phenethyl and naphthylmethyl; Z is $C=O$; B is a bond or a C_1-5 alkylene, C_2-5 alkenylene or C_4-6 alkadienylene group; and Ar is thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, triazolyl, tetrazolyl, tetrahydrofuryl, pyranal, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, piperidyl, piperazinyl, morpholinyl, triazinyl, thiazinyl, benzothienyl, benzofuryl, isobenzofuryl, indolyl, isoindolyl, indazolyl, purinyl, quinolyl, isoquinolyl, naphthyridinyl, quinoxalinyl, chromanyl, indolinyl, isoindolinyl, or chromenyl, unsubstituted or substituted by at least one substituent selected from halogen, hydroxyl, amino, carboxy, C_1-8 alkyl, oxo, C_1-8 alkoxy, phenyl, naphthyl, chlorophenyl, benzyl, phenethyl, naphthylmethyl, (2,3,4- or 3,4,5-)trimethoxybenzyl, amino- C_1-4 alkyl, di- C_1-4 alkylamino- C_1-4 alkyl, hydroxy-substituted C_1-4 alkyl, C_1-5 alkanoyloxy, sterylcarbonyl, 3,4,5-trimethoxystyrylcarbonyl, benzyloxycarbonylamino, benzyloxycarbonylaminomethyl, benzoylamino, 3,4,5-trimethoxybenzoylamino, pyrrolidinylcarbonylmethyl, piperidylcarbonylmethyl, piperazinylcarbonylmethyl and morpholinylcarbonylmethyl are disclosed as anti-ulcer agents in U.S. Patent No. 4,610,983.

The term "loweralkyl" as used herein refers to straight or branched chain alkyl radicals containing from 1 to 8 carbon atoms including but not limited to methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, n-pentyl, 2-methylbutyl, 2,2-dimethylpropyl, n-hexyl, 2-methylpentyl, 2,2-dimethylbutyl, n-heptyl, n-octyl and the like.

The term "cycloalkyl" as used herein refers to an alicyclic ring having 3 to 8 carbon atoms, including, but not limited to cyclopropyl, cyclopentyl, cyclohexyl and the like.

The term "loweralkenyl" or "alkylene group" as used herein refers to a $(CH_2)_y$ radical where y is 1 to 8, such as methylene, ethylene, propylene, tetramethylene and the like.

The term "loweralkynyl" as used herein refers to a lower alkyl radical which contains at least one carbon-carbon triple bond.

The term "alkenylene group" as used herein refers to a C_2-C_8 chain of carbon atoms which contains at least one carbon-carbon double bond, such as vinylene, propenylene, butenylene and the like.

The term "alkadienylene group" as used herein refers to a C_4-C_8 chain of carbon atoms containing at least two carbon-carbon double bonds, such as 1,3-butadienylene and the like.

The term "halogen" as used herein refers to F, Cl, Br or I.

The terms "alkoxy" and "thioalkoxy" as used herein refer to R_8O- and R_8S- respectively, wherein R_8 is a loweralkyl group.

The term "alkoxyalkyl" as used herein refers to an alkoxy group appended to a loweralkyl radical, including, but not limited to methoxymethyl, 2-methoxyethyl and the like.


The term "aryl" or "aryl group" as used herein refers to phenyl, naphthyl, indanyl, fluorenyl, (1,2,3,4)-tetrahydronaphthyl, indenyl or isoindenyl.

The term "substituted aryl" as used herein refers to an aryl group substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH(loweralkyl).

The term "arylalkyl" as used herein refers to an aryl group appended to a loweralkyl radical, including, but not limited to benzyl, phenethyl, naphthylmethyl and the like.

The term "(substituted aryl)alkyl" as used herein refers to a substituted aryl group appended to a loweralkyl radical, including, but not limited to halobenzyl, alkoxynaphthylmethyl and the like.

The term "heterocyclic" or "heterocyclic group" as used herein refers to mono-, bi-, and tricyclic ring systems containing from one to four heteroatoms selected from the group consisting of nitrogen, sulfur, and oxygen; and having from 0 to 7 double bonds. Preferred heterocyclics are: thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, oxatriazolyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolyl, pyrazolidinyl, triazolyl, tetrazolyl, tetrahydrofuryl, pyranal, pyronyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, piperidyl, piperazinyl, morpholinyl, triazinyl, oxazinyl, isoxazinyl, oxathiazinyl, thiazinyl, oxadiazinyl, azepinyl, thiapinyl, thionaphthyl, benzofuryl, isobenzofuryl, indolyl, oxyindolyl, isoindolyl, indazolyl, indolyl, 7-azaindolyl, isoindazolyl, anthranal, benzopyranal, coumarinyl, isocoumarinyl, purinyl, quinolyl, isoquinolyl, naphthyridinyl, cinnolyl, quinazolinyl, pyridopyridyl, benzoxazinyl, diazepinyl, quinoxadinyl, chromenyl, chromanyl, isochromanyl, carbolinyl, xanthenyl, and acridinyl and



25

30


35


40

45

50

55





wherein E is selected from, $(CH_2)_p$, $(CH=CH)_r$, $C(O)CH_2$, $CH_2(O)$, QCH_2 , CH_2Q , $C(O)Q$, and $YC(O)$ wherein Q is S, O or NR_{10} wherein R_{10} is hydrogen, $-(C(O))_r(C_1-C_8 \text{ loweralkyl})$, $-(C(O))_r \text{ cycloalkyl}$, $-(C(O))_r \text{ loweralkenyl}$, $-(C(O))_r \text{ loweralkynyl}$, thioalkoxy, $-(C(O))_r(CH_2)_m \text{ aryl}$, $-(C(O))_r(CH_2)_m \text{ (substituted aryl)}$ wherein substituted aryl is as defined above, $-(CH_2)_m \text{ carboxyl}$, $-(CH_2)_s(C(O))_r \text{ aryl}$, $-(CH_2)_s(C(O))_r \text{ (substituted aryl)}$, $-(CH_2)_m \text{ carboalkoxy}$, $-(CH_2)_m \text{ carboxamide}$, $-(CH_2)_m \text{ carboaryloxy}$, $-C(X)_3$ wherein X is halogen, $-(C(O))_r(CH_2)_s R_{11}$ wherein R_{11} is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, $-(CH_2)_m \text{ aryl}$, $-(CH_2)_m \text{ (substituted aryl)}$ wherein substituted aryl is as defined above, $-(CH_2)_m \text{ heterocyclic}$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as previously defined, $-NR_4OH$ wherein R_4 is hydrogen, $-(C(O))_r \text{ loweralkyl}$, $-(C(O))_r \text{ loweralkenyl}$, $-(C(O))_r \text{ cycloalkyl}$, $-(C(O))_r \text{ loweralkynyl}$, $-(C(O))_r(CH_2)_m \text{ aryl}$, $-(C(O))_r(CH_2)_m \text{ (substituted aryl)}$ wherein substituted aryl is as defined above, $-(CH_2)_m OR_4$ wherein R_4 is as defined above, or $-(CH_2)_m SR_4$ wherein R_4 is as defined above; with the proviso that Y and W are not both C; Y and W are independently selected from S, O, C, CH, CH_2 and NR_{10} with the proviso that Y is not C, CH or CH_2 when W is C, CH or CH_2 ; and wherein R_5 is one, two or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_6 \text{ loweralkyl})$, $(C(O))_r \text{ cycloalkyl}$, decahydronaphthyl, $-(C(O))_r \text{ loweralkenyl}$, $-(C(O))_r \text{ loweralkynyl}$, thioalkoxy, $-(C(O))_r(CH_2)_m \text{ aryl}$, $-(C(O))_r(CH_2)_m \text{ (substituted aryl)}$, $-(CH_2)_m \text{ carboxyl}$, $-(CH_2)_s(C(O))_r \text{ aryl}$, $-(CH_2)_s(C(O))_r \text{ (substituted aryl)}$, $-(CH_2)_m \text{ carboalkoxy}$, $-(CH_2)_m \text{ carboxamide}$, $-(CH_2)_m \text{ carboaryloxy}$, $-C(X)_3$ wherein X is halogen, $-(CO)_r(CH_2)_m R_6$ wherein R_6 is as previously defined, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as previously defined, $-NR_4OH$ wherein R_4 is as previously defined, $-(CH_2)_m OR_4$ wherein R_4 is as defined above, $-(CH_2)_m SR_4$ wherein R_4 is as defined above; m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4.

The symbols used to define the heterocyclic groups in which R_5 is attached to a bond which extends across 2 or 3 rings is meant to indicate that R_5 groups may be bonded to one, two or three of the rings of the heterocyclic ring system.

The term "aryloxy" as used herein refers to R_9O- , wherein R_9 is an aryl or substituted aryl group.

The term "N-protecting group" or "N-protected" as used herein refers to those groups intended to protect the N-terminus of an amino acid or peptide or to protect an amino group against undesirable reactions during synthetic procedures or to prevent the attack of exopeptidases on the compounds or to increase the solubility of the compounds and includes but is not limited to sulfonyl, acyl, acetyl, pivaloyl, t-butyloxycarbonyl (Boc), carbonylbenzyloxy (Cbz), benzoyl or an L- or D-aminoacyl residue, which may itself be N-protected similarly.

The term "O-protecting group" as used herein refers to a substituent which protects hydroxyl groups against undesirable reactions during synthetic procedures and includes but is not limited to substituted methyl ethers, for example methoxymethyl, benzyloxymethyl, 2-methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, benzyl and triphenylmethyl; tetrahydropyranyl ethers; substituted ethyl ethers, for example, 2,2,2-trichloroethyl and t-butyl; silyl ethers, for example, trimethylsilyl, t-butyl dimethylsilyl and t-butyl diphenylsilyl; cyclic acetals and ketals, for example, methylene acetal, acetonide and benzylidene acetal; cyclic ortho esters, for example, methoxymethylene; cyclic carbonates; and cyclic boronates.

The terms "Glu" and "Asp" as used herein refer to glutamic acid and aspartic acid respectively.

The compounds of the invention may be made as shown in Scheme I. Compounds with DL, L, or D configuration may be used in the synthetic schemes.

As illustrated in Scheme I, N-protected omega-benzyl esters of alpha-aminodicarboxylic acids 1 -

(preferably Boc-(gamma-benzyl ester)-D-glutamic acid or Cbz-(gamma-benzyl ester)-D-glutamic acid) are coupled with a primary or secondary amine (preferably dialkylamines, diaryl amines, or alkylaryl amines) using bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOPCl) or a conventional peptide coupling reagent (isobutyl chloroformate (IBCF), phosphorus pentachloride (PCl₅) and the like). The product amide 2 (P is Boc) is deprotected (preferably with hydrochloric acid (HCl) in dioxane) to provide the

hydrochloride salt 3, or 2 is deprotected (P is Cbz) to provide the free amine 4 (preferably using a transfer hydrogenolysis technique). The hydrochloride 3 is coupled with an arylcarboxylic acid (preferably indole-2-carboxylic acid or quinoline-3-carboxylic acid) using 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI) and triethyl amine in dimethylformamide (DMF) as solvent. Other conventional peptide coupling techniques may also be used. Substituents R'CH₂ can be introduced using the appropriate aldehyde (R'CHO) and

amine hydrochloride 3 in a reductive amination reaction (preferably sodium cyanoborohydride or sodium borohydride in methanol or other alcohols as solvents) to provide compound 3a which can be coupled in analogous fashion to 3 to provide 5. Alternatively, the hydrochloride 3 can be directly alkylated with an alkylating agent (R'CH₂X wherein X is Cl, Br, I, OTs and the like) to provide the substituted amine 3a. The product amide 5 is debenzylated utilizing transfer hydrogenation techniques (preferably cyclohexadiene and

palladium on carbon (Pd/C) in ethanol or other alcohols) to provide acid 6. Acid 6 also can be obtained from 5 through standard saponification conditions (aqueous hydroxide in alcohol, THF or dioxane). Alternatively the product 6 is obtained by coupling the amine 4 with active esters of arylcarboxylic acids (preferably indole-2-carboxy-2,4,6-trichlorophenyl ester). The acid 6 and alcohol (R₄OH) are coupled using dicyclohexylcarbodiimide (DCC) or other conventional peptide coupling reagent to provide ester 7. Acid 6 and

secondary amine (NHR₅R₇) likewise are coupled with DCC (or other conventional peptide coupling reagent) to provide product 8. Alternatively the primary amine (R₄NH₂) can be used in place of the secondary amine to provide product 9. Product 9 is also obtained by the direct reaction of benzyl ester 5 and the primary amine (R₄NH₂) at high temperatures and/or pressures.

Reaction of hydrochloride 3 with arylsulfonylchloride in the presence of base (preferably an organic tertiary amine such as triethylamine or N-methylmorpholine) provides product 10. Sulfonamide 10 is debenzylated in a similar fashion to 5 via either transfer hydrogenation conditions or saponification in alcoholic hydroxide. Reaction of hydrochloride salt 3 with a beta-arylacrylic acid (preferably 3-(3-pyridyl)-beta-acrylic acid) utilizing DCC (or other conventional peptide coupling technique) provides benzyl ester 12, which can be deprotected via either transfer hydrogenation conditions or saponification to provide


product acid 13.


In addition, the hydrochloride salt 3 can be coupled, in the presence of base, with acylating agents that are available to one skilled in the art of organic synthesis, including, but not limited to isocyanates, isothiocyanates, acylcyanides, chloroformates and the like. These can be deprotected as previously outlined via hydrogenolysis or saponification to provide ureas, thioureas carbamates and the like.


The N-carbobenzyloxy protected asparagine or glutamine 14 (preferably of the enantiomerically pure R configuration) is converted to its corresponding nitrile 15 by reaction with phosphorus oxychloride and pyridine (alternatively DCC and pyridine). The acid nitrile 15 is coupled with an appropriate amine (preferably dialkyl, diaryl, or alkylarylamine) using DCC (or a conventional coupling reagent) to yield the amide 16. The nitrile 16 is converted to the tetrazole 17 via reaction with sodium azide (NaN₃) in

dimethylformamide (DMF). The tetrazole 17 is deprotected using standard hydrogenation (hydrogen atmosphere over palladium on carbon as catalyst) or transfer hydrogenation conditions to provide amine 18, which is reacted with an active ester of the desired arylcarboxylic acid to provide product 19.

SCHEME I







The following examples will serve to further illustrate preparation of the novel compounds of the invention.

EXAMPLE 1

N-Boc-(γ -benzyl ester)-R-Glu-di-n-pentylamide

To the solution of Boc-(γ -benzyl ester)-R-Glutamic acid (4.0 g, 11.9 mmol) in tetrahydrofuran (THF) (100 mL) was added bis (2-oxo-3-oxazolidinyl)-phosphinic chloride (BOPCl) (3.1 g, 11.9 mmol) and di-n-pentylamine (6 mL, 30 mmol) at room temperature. The reaction mixture was allowed to stir overnight. The solvents were evaporated and the residue dissolved in ethyl acetate (300 mL) and washed with 1N hydrochloric acid (HCl), saturated NaHCO₃ (sodium bicarbonate), H₂O and brine and dried over magnesium sulfate (MgSO₄). After evaporation of solvents the product was subjected to flash chromatography using ethyl acetate (EtOAc)-hexane (2:8) as the eluant. The pure oily product was isolated in 78.1% yield (4.5 g). $[\alpha]_D^{25} = -26.1^\circ$ (c = 1.3, methanol (MeOH)). MS(Cl) m/e 477(m+H)⁺, 421, 381, 363, 325, 299, 282. ¹H NMR(CDCl₃, 300MHz) δ 0.9(m, 6H), 1.3(m, 8H), 1.4(s, 9H), 1.6(m, 4H), 1.7(m, 2H), 2.5(m, 2H), 3.1-3.2(m, 2H), 3.5-4.6(m, 1H), 5.1(s, 2H) 5.4(d, 1H, J = 9 Hz), 7.4(s, 5H).

EXAMPLE 2

(γ -Benzyl ester)-R-Glu-di-n-pentylamide hydrochloride

t-Butyloxycarbonyl derivative of example 1, (1.2 g, 2.5 mmol) was dissolved in 5 mL of 4M HCl in dioxane and the mixture was stirred for 39 min. The solvents were evaporated and the residue was triturated several times with ether and the volatiles were evaporated. The product was filtered and washed several times with diethyl ether. Yield 0.9 g (96%). MS(Cl) m/e 377(m+H)⁺, 359, 269, 241. ¹H NMR-(DMSO-d₆, 300MHz) δ 0.8-0.9(m, 6H), 1.15-1.35(m, 8H), 1.4-1.6(m, 4H), 1.90-2.05(m, 2H), 2.45-2.70(m, 2H), 3.05-3.15(m, 1H), 3.15(m, 1H), 3.30(H₂O), 3.35-3.55(m, 2H), 4.23(m, 1H), 5.1(bs, 2H), 7.37(bs, 5H), 8.2-8.4(bs, 2H).

EXAMPLE 3

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu -di-n-pentylamide

To a solution of hydrochloride of example 2 (7 g, 17 mmol) and N-methylmorpholine (NMM) (3.8 mL, 34.5 mmol) in dimethylformamide (DMF) at 0°C was added indole-2-carboxylic acid (2.75 g, 17 mmol), 1-hydroxybenzotriazole (HOBt) (4.9 g, 36.3 mmol) followed by 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDCI) (3.25 g, 17 mmol). The reaction mixture was allowed to warm to ambient temperature and was stirred overnight. The solvent was concentrated under vacuo and after addition of water, the mixture was extracted successively with 1 N hydrochloric acid, water, saturated solution of sodium bicarbonate and brine. The solution was dried over magnesium sulfate and filtered. Purification was accomplished by crystallization from ethyl acetate and hexane providing 7.55 g (85.6%) of product. $[\alpha]_D^{25} = +24.00^\circ$ (c=0.5, MeOH). mp = 84-86°C. MS(Cl) m/e 520(m+H)⁺, 429, 286, 144. ¹H NMR(CDCl₃, 300MHz) δ 0.85-0.95(m, 6H), 1.2-1.4 (m, 8H), 1.50-1.66(m, 4H), 2.0(m, 1H), 2.20(m, 1H), 2.40-2.65(m, 2H), 3.1(m, 1H), 3.25(m, 1H), 3.48-3.62(m, 2H), 5.05(d, 1H, J = 12 Hz), 5.14 (d, 1H, J = 12 Hz), 5.18(dt, 1H, J = 3.9 Hz), 6.95(d, 1H, J = 2 Hz), 7.15(dt, 1H, J = 1.7.5 Hz), 7.3(m, 2H), 7.35(m, 5H), 7.4(d, 1H, J = 7.5 Hz), 7.65(d, 1H, J = 7.5 Hz), 9.22(bs, 1H). Analysis calculated for C₃₁H₄₁N₃O₄ H₂O: C 71.03, H 7.98, N 8.02; Found: C 71.05, H 7.98, N 8.03.

EXAMPLE 4

N-(2'-Indolylcarbonyl)-R-Glu-N α ,N α -di-n-pentylamide

To a suspension of 0.9 g 10% palladium on carbon (Pd/C) in methanol (10 mL) and cyclohexadiene (3 mL) was added under nitrogen atmosphere (N₂) a solution of benzyl ester of example 3 (1.3g, 2.5 mmol) in

methanol by cannula. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with methanol. The filtrate and washings were combined and concentrated under vacuo. The product was purified by flash chromatography using chloroform (CHCl₃)-methanol and ammonium hydroxide (NH₄OH) 90:10:1 as elutant. Lyophilization provided product in 83.8% yield (0.9 g). mp = 78-79° C. [α]_D = + 2.8° (c = 1.0, MeOH). MS(FAB) m/e 430(m+H)⁺, 273, 245, 226. ¹H NMR(DMSO-d₆, 300MHz) δ 0.8-0.9(m, 6H), 1.12-1.38(m, 8H), 1.39-1.5(m, 4H), 1.56-1.81(m, 2H), 1.85-2.0-(m, 2H), 2.3(t, J=6Hz, 2H), 3.18(m, 1H), 3.54-3.62(m, 2H), 4.85(m, 1H), 7.03(t, J=7Hz, 1H), 7.18(t, J=7Hz, 1H), 7.25(s, 1H), 7.43(d, J=9Hz, 1H), 7.61(d, J=9Hz, 1H), 8.8 (d, J=8Hz, 1H), 11.6(bs, 1H). Analysis calculated for C₂₄H₃₅N₃O₄ H₂O: C 65.77, H 8.18, N 9.59. Found: C 65.99, H 8.01, N 9.76.

EXAMPLE 5

N-(2'-Quinolylcarbonyl)- γ -benzyl ester)-R-Gludi-n-pentylamide

To a solution of hydrochloride of example 2 (0.25g, 0.61 mmol) and N-methylmorpholine (0.14 mL, 1.2 mmol) in DMF (5 mL) at 0° C was added HOBt (0.16 g, 1.2 mmol) and quinoline-2-carboxylic acid (0.11 g, 0.61 mmol) followed by EDCI (0.125 g, 0.65 mmol). The reaction mixture was stirred overnight at room temperature. The solvent was evaporated. After addition of water, the mixture was extracted with ethyl acetate. Combined ethyl acetate extracts were washed with 1.0 N hydrochloric acid, water, a saturated solution of sodium bicarbonate and brine. The solution was dried over magnesium sulfate. The ethyl acetate was evaporated and residue was purified by chromatography using hexane and ethyl acetate as elutant. The product was isolated in 33% yield (0.1 g). MS(Cl) m/e 532(m+H)⁺, 347, 156, 128, 91. ¹H NMR-(CDCl₃, 300MHz) δ 0.85-0.95(m, 6H), 1.2-1.45(m, 8H), 1.6(m, 4H), 2.05(m, 1H), 2.25(m, 1H), 2.46-2.68(m, 2H), 3.1(m, 1H), 3.3(m, 1H), 3.6(m, 2H), 5.12(s, 2H), 5.25(dt, J=3.9Hz, 1H), 7.4(m, 5H), 7.6(dt, J=1.6Hz, 1H), 7.78-(dt, J=1.6Hz, 1H), 7.9(d, J=9Hz, 1H), 8.15(d, J=9Hz, 1H), 8.25(d, J=3Hz, 1H), 8.30(d, J=9Hz, 1H), 9.1-(d, J=9Hz, 1H).

EXAMPLE 6

N-(2'-Quinolylcarbonyl)-R-Glu-N α ,N α -di-n-pentylamide

To a suspension of 0.1 g 10% Pd/C in methanol (3 mL) and cyclohexadiene (1 mL) was added under inert atmosphere (N₂) a solution of benzyl ester of example 5 (0.09 g, 0.170 mmol) in methanol by cannula. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with methanol. The filtrate and washings were combined and concentrated in vacuo. The product was purified by flash chromatography using chloroform-methanol and ammonium hydroxide 90:10:1 as elutant. Lyophilization provided product in 60% yield (0.045 g). MS(Cl) m/e 442-(m+H)⁺, 332, 286, 257, 156, 128. ¹H NMR(CDCl₃, 300MHz) δ 0.82-0.92(m, 6H), 1.2-1.4(m, 8H), 1.50-1.68-(m, 4H), 2.0(m, 1H), 2.2(m, 1H), 2.5(m, 2H), 3.1(m, 1H), 3.25(m, 1H), 3.48-3.70(m, 2H), 5.2(dt, J=3.9Hz, 1H), 7.60-(dt, J=1.6Hz, 1H), 7.75(dt, J=1.6Hz, 1H), 7.85(d, J=6Hz, 1H), 8.18(d, J=9Hz, 1H), 8.25(d, J=6Hz, 1H), 8.31-(d, J=9Hz, 1H), 9.2(d, J=6Hz, 1H). Analysis calculated for C₂₅H₃₅N₃O₄ 0.5 H₂O: C 66.64, H 7.95, N 9.33. Found: C 66.65, H 7.87, N 9.21.

EXAMPLE

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-S-Glu-di-n-pentylamide

To a solution of hydrochloride prepared analogously as in example 2 (0.92 g, 2.23 mmol) and N-methylmorpholine (0.5 mL, 4.5 mmol) in DMF at 0 °C was added indole-2-carboxylic acid (0.36 g, 2.23 mmol), HOBT (0.45 g, 3.35 mmol) followed by EDCI (0.44 g, 2.23 mmol). The reaction mixture was allowed to warm to ambient temperature and was stirred overnight. The solvent was concentrated in vacuo. After addition of ethyl acetate and water, the mixture was extracted with 1.0 N hydrochloric acid, water, a saturated solution of sodium bicarbonate and brine. The solution was dried over magnesium sulfate and filtered. Purification was accomplished by crystallization from ethyl acetate and hexane providing 0.8 g (69.1 %) of product. mp = 83-84 °C. $[\alpha]_D^{25} = -23.55^\circ$ (c=1.0, MeOH). MS(Cl) m/e 520(m+H)⁺. ¹H NMR(CDCl₃, 300MHz) δ 0.8-0.9(m, 6H), 1.2-1.45(m, 8H), 1.52-1.6(m, 4H), 2.0(m, 1H), 2.2(m, 1H), 2.4-2.65(m, 2H), 3.1(m, 1H), 3.25(m, 1H), 3.5-3.68(m, 2H), 5.1(d, J=4Hz, 1H), 5.15(d, J=12Hz, 1H), 5.2(m, 1H), 6.9(d, J=3Hz, 1H), 7.15(t, J=9Hz, 1H), 7.3(m, 1H), 7.35(m, 6H), 7.4(d, J=9Hz, 1H), 7.65(d, J=9Hz, 1H), 9.2(s, 1H).

EXAMPLE 8

N-(2'-Indolylcarbonyl)-S-Glu-N^α, N^α-di-n-pentylamide

To a suspension of 0.5 g 10% Pd/C in methanol (10 mL) and cyclohexadiene (3 mL) was added under inert atmosphere (N₂) solution of benzyl ester of example 7 (0.8 g, 1.55 mmol) in methanol by cannula. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with methanol. The filtrate and washings were combined and concentrated in vacuo. The product was purified by flash chromatography using chloroform-methanol and ammonium hydroxide 90:10:1 as eluant. Lyophilization provided product in 89.5 % yield (0.6 g). mp = 76-74 °C. $[\alpha]_D^{25} = -6.5^\circ$ (c=0.8, MeOH). MS(Cl) m/e 430(m+H)⁺, 273, 245, 144. ¹H NMR(DMSO-d₆, 300MHz) δ 0.8-0.9(m, 6H), 1.12-1.45(m, 8H), 1.48-1.5(m, 4H), 1.53-1.75(m, 2H), 1.80-2.0(m, 2H), 2.3(t, J=6Hz, 2H), 3.0-3.13(m, 2H), 4.85(m, 1H), 7.03(t, J=9Hz, 1H), 7.18(t, J=9Hz, 1H), 7.25(s, 1H), 7.4(d, J=9Hz, 1H), 7.6(d, J=9Hz, 1H), 8.6(d, J=9Hz, 1H), 11.6(s, 1H), 12.2(bs, 1H). Analysis calculated for C₂₄H₃₅N₃O₄ 0.33 H₂O: C 66.19, H 8.25, N 9.65. Found: C 66.39, H 8.23, N 9.65.

EXAMPLE 9

N-(2'-Quinoxalylcarbonyl)-(γ-benzyl ester)-R-Gludi-n-pentylamide

To the stirred solution of hydrochloride of example 2 (0.37 g, 0.9 mmol) and pyridine (0.2 mL, 2 mmol) in anhydrous DMF (2 mL) at 0 °C was added dropwise a solution of 2-quinoxalyl chloride (0.8 g, 0.95 mmol) in DMF (2 mL) over 2 hours. The reaction was monitored by tic using hexane-ethylacetate mixture in 2:1 ratio. Solvents were evaporated and the residue diluted with ethyl acetate and washed successively with 1.0 N hydrochloric acid, water, a solution of saturated bicarbonate and brine. The solution was dried over magnesium sulfate. The product was purified by chromatography (EtOAc/hexane) to provide 0.355 g (74%). MS(Cl) m/e 533(m+H)⁺, 377, 348. ¹H NMR(CDCl₃, 300MHz) δ 0.82-0.98(t, J=6Hz, 6H), 1.23-1.45(m, 8H), 1.5-1.7(m, 4H), 2.05(m, 1H), 2.3(m, 1H), 2.5-2.65(m, 2H), 3.1(m, 1H), 3.3(m, 1H), 3.5-3.72(m, 2H), 5.15(s, 2H), 5.25(dt, J=4.9Hz, 1H), 7.35(m, 5H), 7.85(m, 2H), 8.15(dd, J=1.5, 12Hz, 2H), 8.78(d, J=9Hz, 1H), 9.62(s, 1H).

EXAMPLE 10

N-(2'-Quinoxalylcarbonyl)-R-Glu-N^α, N^α-di-n-pentylamide

To a suspension of 0.08 g 10% Pd/C in absolute ethanol (EtOH, 4 mL) and cyclohexadiene (1 mL) was added under inert atmosphere (N₂) solution of benzyl ester of example 9 (0.08 g, 0.15 mmol) in ethanol via cannula. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with ethanol. The filtrate and washings were combined and concentrated under vacuo. The product was purified by flash chromatography using chloroform-methanol and ammonium hydroxide 90:10:1 as elutant. Lyophilization provided product in 90.0% yield (0.06 g). $[\alpha]_D^{25} = +12.67^\circ$ (c = 0.6, MeOH). MS(Cl) m/e 443(M+H)⁺, 258. ¹H NMR (CDCl₃, 300MHz) δ 0.78-0.98(m, 6H), 1.2-1.4(m, 8H), 1.48-1.65(m, 4H), 1.8(m, 1H), 2.25(m, 1H), 3.05(m, 1H), 3.25(m, 1H), 3.5-3.75(m, 2H), 4.18(m, 2H), 5.3(t, J=9Hz, 1H), 7.8(m, 2H), 8.1(m, 2H), 8.9(d, J=9Hz, 1H), 9.5(s, 1H).

EXAMPLE 11

N-(3'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-di-n-pentylamide

To the stirred solution of hydrochloride of example 2 (1.15 g, 2.8 mmol) in DMF at 0°C and N-methylmorpholine (0.65 mL, 5.6 mmol) was added HOBt (0.75 g, 5.6 mmol) and quinoline-3-carboxylic acid (0.45 g, 2.9 mmol) followed by EDCI (0.57 g, 3.0 mmol). The reaction was allowed to warm to ambient temperature over 2 hours and stirred overnight. Solvent was evaporated and to the residue was added water and ethyl acetate. The organic extracts were washed successively with 1 N hydrochloric acid, water, a solution of saturated sodium bicarbonate and brine. The solution was dried over magnesium sulfate and filtered. The product was purified by chromatography using ethyl acetate and hexane as eluant. Product was obtained in 56.6% yield (0.85 g). MS(Cl) m/e 520(m+H)⁺, 363, 335, 279. ¹H NMR(CDCl₃, 300MHz) δ 0.85-0.95(m, 6H), 1.2-1.42(m, 8H), 1.5-1.7(m, 4H), 1.88-2.02(m, 1H), 2.25(m, 1H), 2.6(m, 2H), 3.1(m, 1H), 3.28(m, 1H), 3.6(m, 2H), 5.1(s, 2H), 5.3(dt, J=3.9Hz, 1H), 7.1(d, J=7.5Hz, 1H), 7.25(m, 2H), 7.3(m, 5H), 7.5(m, 1H), 7.8(d, J=2Hz, 1H), 8.1(m, 1H), 8.9(s, 1H).

EXAMPLE 12

N-(3'-Indolylcarbonyl)-R-Glu-N α ,N α -di-n-pentylamide

To a suspension of 0.3 g 10% Pd/C in absolute ethanol (5 mL) and cyclohexadiene (2 mL) was added under inert atmosphere (N₂) a solution of benzyl ester of example 11 (0.41 g, 0.79 mmol) in ethanol by cannula. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with ethanol. The filtrate and washings were combined and concentrated under vacuo. The product was purified by flash chromatography using chloroform-methanol and ammonium hydroxide 90:10:1 as elutant. Lyophilization provided product in 65% yield (0.22 g). MS(Cl) m/e 430(m+H)⁺, 269, 245. ¹H NMR (DMSO-d₆, 300MHz) δ 0.8-0.9(m, 6H), 1.15-1.32(m, 8H), 1.4-1.72(m, 4H), 1.78-1.9 (m, 2H), 2.20(m, 2H), 3.1-3.4(m, 2H), 3.6(m, 2H), 4.95(m, 1H), 7.12(m, 2H), 7.41(d, J=9Hz, 1H), 7.93(d, J=7Hz, 1H), 8.1(d, J=6Hz, 1H), 8.18(d, J=2Hz, 1H), 11.5(s, 1H), 12.15(bs, 1H). Analysis calculated for C₂₄H₃₅N₃O₄: C 67.10, H 8.21, N 9.78. Found: C 66.97, H 8.30, N 9.75.

EXAMPLE 13

N-(1',2',3',4'-Tetrahydroquinoxaly-2'-carbonyl)-R-Glu-N α ,N α -di-n-pentylamide

To a suspension of 0.3 g 10% Pd/C in absolute ethanol (5 mL) and formic acid (80%) (1 mL) was

added under inert atmosphere (N₂) a solution of benzyl ester of example 9 (0.355 g, 0.67 mmol) and 0.15 g ammonium formate in ethanol via cannula. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with methanol. The filtrate and washings were combined and concentrated in vacuo. The product was purified by flash chromatography using chloroform-methanol and ammonium hydroxide 90:10:1 as elutant. Lyophilization provided product in 67% yield (0.2 g). $[\alpha]_D^{25} = +12.67^\circ$ (c = 0.6, MeOH). MS(Cl) m/e 447(m+H)⁺. ¹H NMR(CDCl₃, 300MHz) δ 0.84-0.96(m, 6H), 1.20-1.38(m, 8H), 1.42-1.70(m, 4H), 1.9-2.02(m, 2H), 2.30(m, 2H), 3.05(m, 1H), 3.18(m, 1H), 3.38-3.6(m, 4H), 3.7(d, J=3Hz, 1H), 3.73(d, J=3Hz, 1H), 4.1(m, 1H), 5.0(dt, J=2.12Hz, 1H), 6.7(d, J=6Hz, 1H), 6.15(d, J=6Hz, 1H), 6.2(m, 2H), 7.85(d, J=9Hz, 1). Analysis calculated for C₂₄H₃₈N₄O₄ 0.9 H₂O: C 62.29, H 8.64, N 12.11. Found: C 62.03, H 8.31, N 12.47.

EXAMPLE 14

N-(3'-Quinolylcarbonyl)-(γ -benzyl ester)-R-Glu-di-n-pentylamide

To a solution of hydrochloride of example 2 (0.41 g, 1 mmol) and N-methylmorpholine (0.22 mL, 2 mmol) in DMF at 0°C was added quinoline-3-carboxylic acid (0.17 g, 1 mmol), HOBt (0.27 g, 2 mmol) followed by EDCI (0.2 g, 1.05 mmol). The reaction mixture was allowed to warm to ambient temperature and then stirred overnight. The solvent was evaporated and water was added to the residue. The mixture was extracted with ethyl acetate. Combined ethyl acetate extracts were washed successively with 1 N hydrochloric acid, water, a saturated solution of sodium bicarbonate, brine and the solution dried over magnesium sulfate then filtered and concentrated in vacuo. Purification by chromatography using ethyl acetate and hexane as elutants provided 0.4 g (75%) of product. MS(Cl) m/e 531(m)⁺, 374, 347, 156, 128, 91. ¹H NMR(CDCl₃, 300MHz) δ 0.85-0.98(m, 6H), 1.2-1.42(m, 8H), 1.5-1.7(m, 4H), 2.02-2.28(m, 2H), 2.50(m, 2H), 3.15(m, 1H), 3.35(m, 1H), 3.5-3.62(m, 2H), 5.15(s, 2H), 5.30(dt, J=3.9Hz, 1H), 7.35(m, 5H), 7.65(t, J=7.5Hz, 1H), 7.85(t, J=7.5Hz, 1H), 7.96(d, J=9Hz, 1H), 8.1(d, J=9Hz, 1H), 8.26(d, J=9Hz, 1H), 8.82(s, 1H), 9.4(d, J=0.5Hz, 1H).

EXAMPLE 15

N-(3'-Quinolylcarbonyl)-R-Glu-N α ,N α -di-n-pentylamide

To a suspension of 0.3 g 10% Pd/C in methanol (10 mL) and cyclohexadiene (3 mL) was added under inert atmosphere (N₂) a solution of benzyl ester of example 14 (0.4 g, 0.75 mmol) in methanol via cannula. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with methanol. The filtrate and washings were combined and concentrated in vacuo. The product was purified by flash chromatography using chloroform-methanol and ammonium hydroxide 90:10:1 as elutant. Lyophilization provided product in 60% yield (0.21g). $[\alpha]_D^{25} = -2.4^\circ$ (c = 0.5, MeOH). MS(FAB) m/e 464(m+Na)⁺, 442(m+H)⁺, 285, 257. ¹H NMR(CDCl₃, 300MHz) δ 0.8-0.95(m, 6H), 1.18-1.4(m, 8H), 1.45-1.7(m, 4H), 2.02(m, 1H), 2.2(m, 1H), 2.54(m, 2H), 3.1(m, 1H), 3.25(m, 1H), 3.5(m, 2H), 5.23(m, 1H), 7.55(t, J=9Hz, 1H), 7.75(t, J=9Hz, 1H), 7.8(d, J=9Hz, 1H), 8.08(d, J=9Hz, 2H), 8.62(d, J=2Hz, 1H), 9.3(d, J=2Hz, 1H). Analysis calculated for C₂₅H₃₅N₃O₄ 0.75 H₂O C 65.98, H 8.09, N 9.23. Found: C 65.92, H 8.21, N 8.86.

EXAMPLE 16

Boc-(β -benzyl ester)-R-Asp-di-n-pentylamide

OuBoc-(β -benzyl ester)-R-Aspartic acid (1.06 g, 3.29 mmol) was stirred at 0 °C in 35 mL of methylene chloride (CH₂Cl₂) under inert atmosphere. Dipentyl amine (7.0 mL) was added to the mixture followed by the addition of BOPCl (1.0 g). The reaction was maintained at 0 °C and when product had formed by tlc the mixture was taken up in ethyl acetate. The organic solvents were washed several times with water then
 5 twice with a 1.0 N hydrochloric acid solution, lastly with a saturated sodium bicarbonate solution. The mixture was dried over magnesium sulfate, filtered and concentrated in vacuo. The residue was purified using a chromatotron and ethyl acetate/hexane mixtures as the elutants to provide 974 mg of product. ¹H NMR(CDCl₃,300MHz) δ 7.35(m,5H), 5.3(bd,J=9Hz,1H), 5.13(d,J=12Hz,1H), 5.08(d,J=12Hz,1H), 4.95-(m,1H), 3.39(m,2H), 3.10-3.30(m,2H), 2.82(dd,J=7,16Hz,1H), 2.63(dd,J=7,16Hz,1H), 1.45-1.65(m,4H), 1.42-
 10 (bs,9H), 1.39(m,8H), 0.89(m,6H).

EXAMPLE 17

(β -Benzyl ester)-R-Asp-di-n-pentylamide hydrochloride

20 Boc-(β -benzyl ester)-R-Asp-di-n-pentyl amide (974 mg) was dissolved in 3.0 mL of a 4.0 N hydrochloric acid solution in dioxane. The reaction was monitored by tlc for disappearance of starting material. When reaction was complete the mixture was diluted with diethyl ether and hexane and volatiles removed in vacuo. This process was repeated three times leaving a thick clear oil which was used for subsequent reactions.

EXAMPLE 18

N-(2'-Indolylcarbonyl)- β -benzyl ester)-R-Asp-di-n-pentylamide

Hydrochloride salt of example 17 (270 mg, 0.67 mmol), indole 2-carboxylic acid (129 mg, 0.80 mmol),
 35 HOBt (100 mg, 0.74 mmol) and EDCI (165 mg, 0.86 mmol) were suspended in 8 mL of anhydrous dimethylformamide at 0 °C. Triethyl amine (TEA) (200 μ L) was added and the reaction mixture stirred under nitrogen allowing warming to ambient temperature overnight. The mixture was taken up in ethyl acetate and washed with water, 1.0 N hydrochloric acid solution, and saturated sodium bicarbonate solution. The organic solution was dried over magnesium sulfate, filtered and the residue purified by chromatography utilizing
 40 chromatotron and ethyl acetate/hexane elutant mixtures to provide 223 mg of product. mp = 94-5 °C. [α]_D = +44.8° (c=0.125, MeOH). MS(Cl) m/e 506(m+H)⁺, 407, 363, 346, 321. ¹H NMR(CDCl₃,300MHz) δ 9.18-(bs,1H), 7.63(bd,J=7.5Hz,1H), 7.42(bd,J=7.5Hz,1H), 7.32(m,7H), 7.14(dt,J=1.5,7.5Hz,1H), 6.85(s,1H), 5.48-(dt,J=6.9Hz,1H), 5.12(m,2H), 3.15-3.50(m,4H), 2.92(dd,J=6,15Hz,1H), 2.29(dd,J=6,15Hz,1H), 1.45-1.60-(m,4H), 1.2-1.35(m,8H), 0.88(m,6H). C,H,N Analysis calculated for C₃₀H₃₉N₃O₄ 0.3 EtOAc: C 70.43, H 7.84, N 7.90; Found C 70.31, H 8.08, N 7.67.

EXAMPLE 19

N-(2'-Indolylcarbonyl)-R-Asp-N α ,N α -di-n-pentylamide

55 To a suspension of 97 mg of 10% Pd on carbon in ethanol under nitrogen atmosphere was added 100 μ L of cyclohexadiene. To this mixture was then added 112 mg of the benzyl ester of example 18 and 400 μ L of cyclohexadiene. The reaction was monitored by tlc for the disappearance of starting material. When reaction was complete the mixture was filtered through celite and the volatiles removed in vacuo. The

residue was lyophilized to provide product. mp = 183-5° C. $[\alpha]_D^{25} = +27.1^\circ$ (c=0.17, acetone). MS(FAB) m/e 438(m+Na)⁺, 416(m+H)⁺, 373, 317, 273. ¹H NMR(DMSO-d₆, 300MHz) δ 11.63(s, 1H), 8.87(d, J=9Hz, 1H), 7.60(d, J=7.5Hz, 1H), 7.42(d, J=7.5Hz, 1H), 7.24(d, J=1.5Hz, 1H), 7.18(dt, J=1.5, 7.5Hz, 1H), 7.02(t, J=7.5Hz, 1H), 5.25(m, 1H), 3.2-3.5(bs, H₂O), 2.84(dd, J=8, 16Hz, 1H), 1.58(bs, 2H), 1.43(m, 2H), 1.15-1.30(m, 8H), 0.85(t, J=7.5Hz, 3H), 0.77(t, J=7.5Hz, 3H). C, H, N Analysis calculated for C₂₃H₃₃N₃O₄ · 0.4 H₂O: C 65.33, H 8.06, N 9.94; Found C 65.32, H 8.16, N 9.72.

EXAMPLE 20

N-(2'-Quinolylcarbonyl)-(β-benzyl ester)-R-Asp-di-n-pentylamide

(β-Benzyl ester)-R-Asp-dipentyl amide hydrochloride (105 mg, 0.26 mmol), quinoline-2-carboxylic acid (57 mg, 0.33 mmol), HOBT (36 mg) and EDCI (55 mg, 0.28 mmol) were suspended in 5 mL of dimethyl formamide stirring at 0° C under nitrogen. To this mixture was added 400 μL of triethyl amine and the reaction mixture was allowed to stir overnight while warming to ambient temperature. The reaction mixture was taken up in ethyl acetate and washed with one portion of water, two portions of 1.0 N hydrochloric acid and a saturated sodium bicarbonate solution. The organics were dried over magnesium sulfate, filtered and the volatiles removed in vacuo. The residue was purified by chromatography to provide 99.5 mg of product. MS(EI) m/e 517(m)⁺, 362, 333, 156, 128, 91. ¹H NMR(CDCl₃, 300MHz) δ 8.87(d, J=9.7Hz, 1H), 8.32(d, J=7.5Hz, 1H), 8.25(d, J=9Hz, 1H), 8.12(bd, J=9Hz, 1H), 7.87(d, J=7.5Hz, 1H), 7.78(dt, J=1.5, 9Hz, 1H), 7.62(dt, J=1.5, 9Hz, 1H), 7.25-7.35(m, 5H), 5.55(m, 1H), 5.17(d, J=12Hz, 1H), 5.10(d, J=12Hz, 1H), 3.4(m, 3H), 3.25(m, 1H), 3.10(dd, J=7.5, 16Hz, 1H), 2.84(dd, J=6, 16Hz, 1H), 1.55(m, 4H), 1.20-1.35(m, 8H), 0.88(t, J=7Hz, 3H), 0.80(t, J=7Hz, 3H).

EXAMPLE 21

N-(2'-Quinolylcarbonyl)-R-Asp-N^α, N^α-di-n-pentylamide

The β-Benzyl ester of example 20 (93 mg, 0.18 mmol) in 2 mL of ethanol were added to 87 mg of 10% palladium on carbon stirring in 4 mL of ethanol under a nitrogen atmosphere. Cyclohexadiene (500 μL) was added and the reaction was monitored for disappearance of starting material. When this had occurred the reaction mixture was filtered through celite and the residue concentrated and then lyophilized from water to provide product (56 mg). mp = 91-3° C. $[\alpha]_D^{25} = +1.25^\circ$ (c=.08, MeOH). MS(FAB) m/e 472(m+2Na-H)⁺, 450(m+Na)⁺, 428(m+H)⁺. ¹H NMR(CDCl₃, 300MHz) δ 9.05(d, J=7.5Hz, 1H), 8.28(d, J=9Hz, 1H), 8.22(d, J=9Hz, 1H), 8.12(bd, J=7.5Hz, 1H), 7.83(bd, J=7.5Hz, 1H), 7.75(bt, J=7.5Hz, 1H), 7.61(bt, J=7.5Hz, 1H), 5.50(dt, J=6.9Hz, 1H), 3.35-3.50(m, 3H), 3.25(m, 1H), 3.04(dd, J=6, 16Hz, 1H), 2.85(dd, J=5, 16Hz, 1H), 2.5-(bs, 1H), 1.60(m, 4H), 1.25(m, 8H), 0.88(t, J=7.5Hz, 3H), 0.80(t, J=7.5Hz, 3H). C, H, N Analysis calculated for C₂₄H₃₃N₃O₄ · 0.5 H₂O: C 66.02, H 7.85, N 9.63; Found C 66.18, H 7.81, N 9.49.

EXAMPLE 22

N-(2'-Indolylcarbonyl)-R-Gln-N^α, N^α-di-n-pentylamide

Ammonia was bubbled into a solution of N-(2'-indolylcarbonyl)-(γ-benzyl ester)-R-Glu-di-n-pentyl amide (50 mg) in methanol at -78° C and the reaction vessel sealed. Stirring continued over 48 hrs with warming to ambient temperature. The volatiles were evaporated in vacuo and product isolated as a white solid (37

mg). mp = 71-4° C. $[\alpha]_D = -2.8^\circ$ (c=0.5, acetone). MS(FAB) m/e 429(m+H)⁺, 396, 272, 244. ¹H NMR-(DMSO-d₆, 300MHz) δ 11.6(bs, 1H), 8.57(bd, J=7.5Hz, 1H), 7.61(d, J=7.5Hz, 1H), 7.42(d, J=7.5Hz, 1H), 7.30-(m, 2H), 7.17(bt, J=7.5Hz, 1H), 7.03(bt, J=7.5Hz, 1H), 6.82(bs, 1H), 4.84(m, 1H), 3.35(bm, H₂O), 3.08(m, 1H), 2.20(m, 2H), 1.75-1.90(m, 2H), 1.70(m, 1H), 1.6(m, 1H), 1.45(m, 2H), 1.2-1.33(m, 8H), 0.86(m, 6H). C, H, N Analysis calculated for C₂₄H₃₆N₄O₃ · 0.2 H₂O: C 66.68, H 8.49, N 12.97; Found C 66.64, H 8.43, N 12.91.

EXAMPLE 23

N-(2'-Indolylcarbonyl)-(γ-methyl ester)-R-Glu-di-n-pentylamide

To a solution of N-(2'-Indolylcarbonyl)-R-Glu-di-n-pentylamide (38 mg) stirring in ethyl acetate at ambient temperature was added an excess of diazomethane solution in ether. The excess diazomethane was destroyed by acetic acid (HOAc) addition and the product was isolated by concentration and filtration through a plug of silica gel providing 28 mg of methyl ester. mp = 93-4° C. $[\alpha]_D = +11.1^\circ$ (c=0.045, MeOH). MS(Cl) m/e 444(m+H)⁺, 287, 269, 158. ¹H NMR(CDCl₃, 300MHz) δ 9.12(bs, 1H), 7.66-(d, J=7.5Hz, 1H), 7.42(bd, J=7.5Hz, 1H), 7.22-7.32(m, 2H), 7.13(dt, J=1.5, 7.5Hz, 1H), 6.95(m, 1H), 5.25-(dt, J=4.9Hz, 1H), 3.68(s, 3H), 3.58(m, 2H), 3.27(m, 1H), 3.10(m, 1H), 2.40-2.60(m, 2H), 2.2(m, 1H), 1.9(m, 1H), 1.5-1.7(m, 4H), 1.25-1.4(m, 8H), 0.91(m, 6H). C, H, N Analysis calculated for C₂₅H₃₇N₃O₄ · 0.5 H₂O: C 66.33, H 8.47, N 9.28. Found C 66.24, H 8.22, N 9.18.

EXAMPLE 24

N-Boc-(γ-benzyl ester)-R-Glu-benzylamide

To a solution of Boc-(γ-benzyl ester)-R-glutamic acid (258 mg) and BOPCI (250 mg) in methylene chloride (7 mL) at 0° C and under nitrogen atmosphere was added benzyl amine (90 μL). The reaction mixture was allowed to stir overnight and reach ambient temperatures. The mixture was taken up in ethyl acetate and extracted with two portions of water, two portions of 0.5 N hydrochloric acid, and one of saturated aqueous sodium bicarbonate solution. The solution was dried over magnesium sulfate and filtered. The solution was concentrated and chromatography of the residue provided 65 mg of product. MS(Cl) m/e 427(m+H)⁺, 399, 371. ¹H NMR(CDCl₃, 300 MHz) δ 7.23-7.46(m, 10H), 6.50(bs, 1H), 5.23(bs, 1H), 5.12(s, 2H), 4.43(bd, J=6Hz, 2H), 4.18(m, 1H), 2.4-2.6(m, 2H), 2.18(m, 1H), 1.96(m, 1H), 1.41(bs, 9H).

EXAMPLE 25

N-(3'-Quinolylcarbonyl)-(γ-benzyl ester)-R-Glu-benzylamide

Boc-(γ-benzyl ester)-R-Glu-di-n-pentyl amide (58 mg) was treated with 4.0 mL of 4.5 M hydrochloric acid in dioxane at 0° C until starting material was consumed by tic analysis. The volatiles were evaporated in vacuo and the residue was stirred with 3-quinolinecarboxylic acid (68 mg), EDCI (74 mg), and HOBT (22 mg) in 4 mL of anhydrous DMF under nitrogen at 0° C. To this reaction was added triethyl amine (75 μL) and the reaction was allowed to warm to ambient temperature overnight. The mixture was taken up in ethyl acetate and extracted with two portions of water, one portion of 1.0 N hydrochloric acid and once with saturated sodium bicarbonate solution. The organic solution was dried over magnesium sulfate and filtered. Removal of the volatiles in vacuo and trituration of the residue with ethyl acetate provided 44 mg of product. mp = 160-62° C. $[\alpha]_D = -15.0^\circ$ (c=.08, acetone). MS(Cl) m/e 482(m+H)⁺, 374, 347. ¹H NMR-

(CDCl₃, 300MHz) δ 9.25(m, 1H), 8.83(m, 1H), 8.08(m, 2H), 7.89(m, 1H), 7.71(m, 1H), 7.4(m, 10H), 4.87(bs, 6H), 4.7(m, 1H), 2.58(bt, J = 7.5Hz, 2H), 2.30(m, 1H), 2.20(m, 1H). C, H, N Analysis calculated for C₂₉H₂₇N₃O₄ 0.833. H₂O: C 70.13, H 5.82, N 8.46; Found C 70.06, H 5.46, N 8.62.

5

EXAMPLE 26

10

N-Boc-(γ -benzyl ester)-R-Glu-dibenzylamide

To a solution of Boc-(γ -benzyl ester)-R-Glutamic acid (715 mg) and BOPCl (750 mg) stirring in 15 mL of methylene chloride at 0° C was added dibenzyl amine (800 mg). The reaction mixture was allowed to stir overnight and warm to ambient temperature. The reaction mixture was taken up in ethyl acetate and extracted twice with 0.5 N hydrochloric acid solution and then with aqueous saturated sodium bicarbonate solution. The solution was dried over magnesium sulfate and filtered. The mixture was concentrated in vacuo and the residue purified by chromatography utilizing ethyl acetate and hexane as the elutants to provide 571 mg of product. MS(Cl) m/e 516(m)⁺, 460, 443, 416, 369, 325, 292, 236. ¹H NMR- (CDCl₃, 300MHz) δ 7.35(m, 11H), 7.18(m, 4H), 5.45(bd, J = 9Hz, 1H), 5.1(m, 2H), 4.7-4.9(m, 3H), 4.45-(d, J = 16Hz, 1H), 4.3(d, J = 16Hz, 1H), 2.45(m, 2H), 2.10(m, 1H), 1.82(m, 1H), 1.43(bs, 9H).

25

EXAMPLE 27(γ -Benzyl ester)-R-Glu-dibenzylamide hydrochloride

30

Boc-(γ -benzyl ester)-R-Glu-dibenzylamide (571 mg) was treated with 7.0 mL of a 4.5 M solution of hydrochloric acid in dioxane at -20° C under nitrogen atmosphere. The reaction was stirred 3 hours at which time tlc indicated no more starting material. The mixture was concentrated in vacuo and the residue triturated with diethyl ether. The volatiles again were evaporated in vacuo and diethyl ether added to the residue. Filtration provided the product as a white solid (475 mg).

40

EXAMPLE 28N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-dibenzylamide

45

EDCl (129 mg), HOBt (10 mg), 2-indolecarboxylic acid (117 mg) and hydrochloride salt of example 27 (236 mg) were suspended in 5 mL of anhydrous DMF under nitrogen at -20° C. To this mixture was added triethyl amine 170 μ l and reaction mixture stirred overnight allowing the reaction to warm to ambient temperature. The reaction mixture was taken up in ethyl acetate and washed with three volumes of water, once with 1.0 N hydrochloric acid and once with saturated aqueous sodium bicarbonate solution. The solution was dried over magnesium sulfate and filtered. Concentration of the solution in vacuo and chromatography of the residue provided 240 mg of product. Crystallization of a small sample from ethyl acetate/hexane provided an analytically pure sample. mp = 94-6° C. [α]_D = +4.5° (c = 0.11, MeOH). MS- (FAB) m/e 560(m+H)⁺, 363, 335, 234, ¹H NMR(CDCl₃, 300MHz) δ 9.05(bs, 1H), 7.68(d, J = 7.5Hz, 1H), 7.29-7.45(m, 14H), 7.20(m, 5H), 7.0(m, 1H), 5.38(dt, J = 3.9Hz, 1H), 5.10(d, J = 15Hz, 1H), 5.05(d, J = 15Hz, 1H), 4.95-(d, J = 15Hz, 1H), 4.82(d, J = 15Hz, 1H), 4.45(d, J = 15Hz, 1H), 4.22(d, J = 15Hz, 1H), 2.40-2.60(m, 2H), 2.25(m, 1H), 2.03(m, 1H). C, H, N analysis calculated for C₃₅H₃₃N₃O₄: C 75.11, H 5.95, N 7.51; Found C 75.63, H 6.00, N 7.55.

EXAMPLE 29N-(2'-Indolylcarbonyl)-R-Glu-N α , N α -dibenzylamide

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-dibenzylamide (210 mg) was stirred in 3 mL of methanol and 3.0 mL of 1.0 N sodium hydroxide (NaOH) solution was added at room temperature. The reaction was stirred until tlc analysis indicated the complete disappearance of starting material. The mixture was then concentrated in vacuo and the residue taken up in ethyl acetate and partitioned with saturated sodium bicarbonate solution. The aqueous solution was separated and acidified with hydrochloric acid to pH 2.0. This solution was extracted three times with equal portions of ethyl acetate and these extracts were combined and dried over magnesium sulfate. Filtration and evaporation of the volatiles in vacuo provided a residue which when triturated with hexane gave product. mp = 111-3 °C. $[\alpha]_D = -5.4^\circ$ (c=0.13, acetone). MS(FAB) m/e 470(m+H)⁺, 273, 234. ¹H NMR(DMSO-d₆, 300 MHz) δ 11.65(bs, 1H), 8.75(d, J=7.5Hz, 1H), 7.63-(d, J=7.5Hz, 1H), 7.15-7.45(m, 14H), 7.05(t, J=7.5Hz, 1H), 5.10(m, 1H), 4.90(d, J=16Hz, 1H), 4.71-(d, J=15Hz, 1H), 4.50(d, J=16Hz, 1H), 4.17(d, J=15Hz, 1H), 2.37(bt, J=7Hz, 2H), 2.0(m, 2H). C,H,N Analysis calculated for C₂₈H₂₇N₃O₄ 0.4 H₂O: C 70.53, H 5.88, N 8.82; Found C 70.49, H 5.79, N 8.79.

EXAMPLE 30Methyl N-Boc-(γ -benzyl ester)-R-Glutamyl-R-phenylglycinate

N-Boc-(γ -benzyl ester)-R-Glutamic acid (692 mg), EDCI (583 mg), HOBt (220 mg), and R-phenylglycine methyl ester hydrochloride (384 mg) were suspended in 15 mL of anhydrous DMF at -20 °C under nitrogen atmosphere. Triethylamine (700 μ L) was added and the reaction mixture was allowed to stir overnight with warming to ambient temperature. The mixture was taken up in a large volume of ethyl acetate and extracted with water, 1.0 N solution of hydrochloric acid, and saturated sodium bicarbonate solution. The solution was dried over magnesium sulfate and then filtered. Evaporation of the volatiles in vacuo and chromatography of the residue provided 832 mg of product. MS(FAB) m/e 485(m+H)⁺, 429, 385, 367, 277, ¹H NMR-(CDCl₃, 300MHz) δ 7.36(brm, 11H), 5.54(d, J=7.5Hz, 1H), 5.23(m, 1H), 5.15(m, 2H), 4.26(m, 1H), 3.72(s, 3H), 2.54-(dt, J=3, 7.5Hz, 2H), 2.18(m, 1H), 1.98(m, 1H), 1.42(s, 9H).

EXAMPLE 31Methyl N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glutamyl-R-phenylglycinate

Boc-(γ -benzyl ester)-R-Glu-R-phenylglycine methyl ester (832 mg) was treated with 6.0 mL of 4.5 M hydrochloric acid in dioxane at 0 °C. When tlc analysis of the reaction mixture indicated complete consumption of the starting material the solvents were removed in vacuo. Several times ether and hexane were added to the residue and solvents evaporated in vacuo. The resulting oily product was used without further purification. Hydrochloride salt (260 mg), 2-indolecarboxylic acid (121 mg), EDCI (163 mg), and HOBt (96 mg) were suspended in 7.0 mL anhydrous DMF at -10 °C and triethyl amine (210 μ L) was added. The reaction mixture was stirred and warmed to ambient temperature overnight. The mixture was taken up in a large volume of ethyl acetate and extracted with two portions of water, two portions of 1.0 N hydrochloric acid solution, and saturated sodium bicarbonate solution. The solution was dried over magnesium sulfate and filtered and the volatiles evaporated in vacuo. Chromatography of the residue utilizing ethyl acetate and hexane mixtures as the elutants provided 262 mg of product. mp = 177-8 °C. $[\alpha]_D = -71.9^\circ$ (c=0.32, DMSO-MeOH (1/1)). MS(Cl) m/e 528(m+H)⁺, 436, 420, 395, 363, 294, 277. ¹H NMR-

(CDCl₃, 300MHz) δ 9.25(bs, 1H), 7.75(m, 1H), 7.67(d, J = 7.5Hz, 1H), 7.26-7.40(m, 13H), 7.14(t, J = 7.5Hz, 1H), 6.94(s, 1H), 5.55(d, J = 7.5Hz, 1H), 5.12(m, 2H), 4.88(m, 1H), 3.74(s, 3H), 2.65(m, 2H), 2.25(m, 2H). C,H,N Analysis calculated for C₃₀H₂₉N₃O₆: C 68.28, H 5.54, N 7.97; Found: C 68.29, H 5.66, N 8.08.

EXAMPLE 32

Methyl N-(3'-Quinolylcarbonyl)-(γ -benzyl ester)-R-Glutamyl-R-phenylglycinate

Hydrochloride salt used in example 31 (373 mg), 3-quinolinecarboxylic acid (168 mg), EDCI (210 mg) and HOBt (96 mg) were suspended in 9.0 mL of anhydrous DMF stirring at -10 °C under nitrogen atmosphere. Triethyl amine (280 μ L) was added and the resulting mixture stirred overnight warming to ambient temperature. The mixture was taken up in a large volume of ethyl acetate and extracted twice with portions of water, twice with 1.0 N hydrochloric acid solutions, and once with saturated sodium bicarbonate. The solution was then dried over magnesium sulfate and filtered. Evaporation of the volatiles and chromatography of the residue utilizing ethyl acetate and hexane mixtures as the elutants provided 367 mg of product. mp = 144-6 °C. $[\alpha]_D^{25} = +69.56^\circ$ (c = 0.11, MeOH). MS(Cl) m/e 540(m + H)⁺, 432, 375, 347, 294. ¹H NMR(CDCl₃, 300MHz) δ 9.32(d, J = 3Hz, 1H), 8.58(m, 1H), 8.15(d, J = 7.5Hz, 1H), 7.9(d, J = 7.5Hz, 1H), 7.83(m, 1H), 7.60(m, 3H), 7.34(m, 10H), 5.55(d, J = 7Hz, 1H), 5.18(m, 2H), 4.83(m, 1H), 3.73(s, 3H), 2.60-2.85-(m, 2H), 2.20-2.40(m, 2H). C,H,N Analysis calculated for C₃₁H₂₉N₃O₆: C 68.99, H 5.42, N 7.79; Found: C 68.88, H 5.45, N 7.78.

EXAMPLE 33

N-(6'-Quinolylcarbonyl)-(γ -benzyl ester)-R-Glu-di-n-pentyl amide

Hydrochloride salt of example 2 (190 mg), EDCI (138 mg), HOBt (86 mg), and 6-quinolinecarboxylic acid (99.6 mg) were suspended in 6.0 mL of anhydrous DMF at 0 °C under a nitrogen atmosphere. Triethyl amine (165 μ L) was added and the reaction mixture stirred overnight with warming to ambient temperature. The reaction mixture was poured into a large volume of ethyl acetate and this solution was extracted twice with water, twice with 1.0 N hydrochloric acid solution, and once with saturated sodium bicarbonate solution. The solution was dried over magnesium sulfate and filtered. Evaporation of the volatiles in vacuo and chromatography of the residue provided 245 mg of product. $[\alpha]_D^{25} = +16.9^\circ$ (c = 0.61, MeOH). MS(Cl) m/e 532(m + H)⁺, 424, 398, 375, 347, 264. ¹H NMR(CDCl₃, 300MHz) δ 8.97(dd, J = 1.5, 5Hz, 1H), 8.30(s, 1H), 8.18-(dd, J = 1.5, 9Hz, 1H), 8.10(bs, 2H), 7.71(bd, J = 8Hz, 1H), 7.44(m, 1H), 7.30-7.35(m, 5H), 5.25(dt, J = 3.5, 9Hz, 1H), 5.14(bs, 2H), 3.60(m, 2H), 3.30(m, 1H), 3.12(dt, J = 7.5, 14Hz, 1H), 2.45-2.70(m, 2H), 2.25(m, 1H), 2.04(m, 1H), 1.68(m, 2H), 1.55(m, 2H), 1.2-1.4(m, 8H), 0.92(m, 6H).

EXAMPLE 34

N-(6'-Quinolylcarbonyl)-R-Glu-N α , N α -di-n-pentylamide

To a suspension of 249 mg of palladium on carbon (10%) in 5 mL of ethanol under nitrogen atmosphere was added 100 μ L of cyclohexadiene, followed by 154 mg of benzyl ester of example 33 in 5 mL of ethanol. To this reaction mixture was added an additional 400 μ L of cyclohexadiene and the reaction was monitored for consumption of starting material. The reaction mixture was filtered through celite and the volatiles removed in vacuo. Crystallization from ethyl acetate and hexane gave 114 mg of product. mp -

146-8° C. $[\alpha]_D = -18.9^\circ$ (c = .095, acetone) MS(Cl) m/e 442(m+H)⁺, 398, 323, 304, 294, 257, 173, 158. ¹H NMR(CDCl₃, 300MHz) δ 8.94(dd, J = 1.5, 4.5Hz, 1H), 8.27(d, J = 1.5Hz, 1H), 8.10-8.18(m, 2H), 8.02(d, J = 9Hz, 1H), 7.70(d, J = 7.5Hz, 1H), 7.42(dd, J = 4.5, 8.5Hz, 1H), 5.24(m, 1H), 3.45-3.65(m, 2H), 3.2-3.35(m, 1H), 3.05-3.17-(m, 1H), 2.55(m, 2H), 2.23(m, 1H), 2.05(m, 1H), 1.65(m, 2H), 1.55(m, 2H), 1.25-1.40(m, 8H), 0.90(m, 6H). C, H, N Analysis calculated for C₂₅H₃₅N₃O₄ 0.33 H₂O: C 67.08, H 8.04, N 9.39; Found: C 67.10, H 7.96, N 9.34.

EXAMPLE 35

Methyl N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-Glutamyl S-phenylglycinate

15 N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-glutamic acid (2.13 gram) and methyl-S-phenylglycinate hydrochloride were reacted in a fashion similar to that in example 21 to provide 1.00 grams (33% yield) of product after chromatography. mp = 91-92° C. $[\alpha]_D = +54.6^\circ$ (c = 0.52, acetone). MS(FAB) m/e 485-(m+H)⁺, 429, 385, 367, 350, 345, 325, 277. ¹H NMR(CDCl₃, 300MHz) δ 7.33(m, 10H), 7.23(m, 1H), 5.53-(d, J = 7.5Hz, 1H), 5.25(m, 1H), 5.13(s, 2H), 4.25(m, 1H), 3.72(s, 3H), 2.52(m, 1H), 2.40(m, 1H), 2.14(m, 1H), 1.94-(m, 1H), 1.42(bs, 9H).

EXAMPLE 36

Methyl N-(2'-Indolylcarbonyl)-(γ-benzylester)-R-Glutamyl-S-phenylglycinate

30 Boc-(γ-benzyl ester)-R-Glu-S-phenylglycine methyl ester (362 mg) was treated with 8.0 mL of 4.5 M hydrochloric acid in dioxane at -10° C under nitrogen atmosphere. Reaction progress was monitored by tlc and when starting material was consumed the solvents were removed in vacuo. EDCI (212 mg), HOBt (100 mg), and indole-2-carboxylic acid (133 mg) were added to this residue and the contents dissolved in 15 mL of anhydrous DMF at 0° C. Triethyl amine (260 μL) was added and the reaction was allowed to stir overnight under nitrogen warming to ambient temperature. The reaction mixture was poured into ethyl acetate and extracted twice with portions of water, twice with portions of 1.0 N HCl solution and once with saturated sodium bicarbonate solution. The solution was dried over magnesium sulfate, filtered, and concentrated in vacuo. Chromatography of the residue provided 230 mg of the desired product. mp = 179-81° C. $[\alpha]_D = +49.4^\circ$ (c = 0.34, MeOH). MS(Cl) m/e 528(m+H)⁺, 437, 420, 395, 294. ¹H NMR-(CDCl₃, 300MHz) δ 9.38(bs, 1H), 7.8(d, J = 7.5Hz, 1H), 7.65(d, J = 7.5Hz, 1H), 7.3-7.45(m, 13H), 7.14-(dt, J = 1.5, 7.5Hz, 1H), 6.98(m, 1H), 5.58(d, J = 7.5Hz, 1H), 5.1(s, 2H), 4.92(m, 1H), 3.71(s, 3H), 2.62(m, 1H), 2.41-(m, 1H), 2.15-2.32(m, 2H). C, H, N Analysis calculated for C₃₀H₂₉N₃O₆: C 68.28, H 5.54, N 7.97; Found: C 68.38, H 5.63, N 7.74.

EXAMPLE 37

Methyl N-(2'-Indolylcarbonyl)-R-Glutamyl-S-phenylglycinate

50 To a suspension of 81 mg of palladium (10%) on carbon in 2 mL of ethanol under nitrogen atmosphere were added cyclohexadiene (100 μL) followed by a solution of benzyl ester of example 36 in 2.0 mL of ethanol. An additional 400 μL of cyclohexadiene was added and the reaction stirred at room temperature until starting material was consumed by tlc analysis. The reaction mixture was filtered through celite and the filtrate concentrated in vacuo. The residue was lyophilized to provide 36 mg of final product. mp = 133-9° C. $[\alpha]_D = +18.4^\circ$ (c = 0.065, acetone). MS(FAB) m/e 438 (m+H)⁺, 391, 273, 245. ¹H NMR-

(DMSO_{d6}, 300MHz) δ 11.63(bs, 1H), 9.43(bs, 1H), 8.92(d, J=7.5Hz, 1H), 7.61(d, J=7.5Hz, 1H), 7.40(m, 7H), 7.23-(bs, 1H), 7.18(t, J=7.5Hz, 1H), 7.03(t, J=7.5Hz, 1H), 5.47(m, 1H), 4.55(m, 1H), 3.62(s, 3H), 3.25-3.50(H₂O), 2.21-(bt, J=7.5Hz, 2H), 1.93(m, 2H). C, H, N Analysis calculated for C₂₃H₂₃N₃O₆ 1.33 H₂O: C 59.85, H 5.56, N 9.11; Found C 59.76, H 5.16, N 8.74.

EXAMPLE 38

N-(2'-Indolylcarbonyl)-R-Glu-N α -(2'-biphenylamide)

To a solution of Boc(γ -benzyl ester)-R-Glutamic acid (337 mg, 1.0 mmol) and 2-aminobiphenyl (423 mg, 2.5 mmol) in THF (15 mL) at 4° C were added TEA (140 mL, 1.0 mmol) and BOPCI (254 mg, 1.0 mmol) and the reaction was allowed to warm to room temperature overnight. The reaction mixture was poured into EtOAc and extracted with portions of 1 M phosphoric acid (H₃PO₄); 1 M Na₂CO₃ (sodium carbonate); H₂O then dried over MgSO₄. Evaporation of the solvent gave a pink oil. Flash chromatography on silica gel eluted with 8:1 hexanes-ethyl acetate gave: 246 mg, (0.51 mmol, 51% yield). R_f = 0.46 (2:1 hexanes-EtOAc); 0.24 (4:1 hexanes-EtOAc). The material above was treated with HCl-dioxane (1.5 mL, 6 mmol) under an N₂ atmosphere for 4 hours at room temperature. The reaction was evaporated in vacuo and then placed under high vacuum overnight. R_f = 0.63 (100:60:18:33 EtOAc-pyridine-HOAc-H₂O). The HCl salt (0.51 mmol), 2-indole carboxylic acid (97mg, 0.60 mmol), HOBT (81 mg, 0.60 mmol) and TEA (167 μ L, 1.2 mmol) were dissolved in CH₂Cl₂ (8 mL) and treated with EDCI (115 mg, 0.66 mmol) at room temperature overnight. The solvent was evaporated in vacuo and the residue was dissolved in EtOAc and extracted with successive portions of 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), H₂O-brine, then dried over MgSO₄, filtered and evaporated. The oily residue was crystallized from give 209 mg, (0.394 mmol, 77%, R_f = 0.23 (2/1 hexane-EtOAc)). The benzyl ester (190 mg, 0.36 mmol) in MeOH (10mL) and acetic acid (HOAc, 5 mL) was treated with 10% Pd/C (prewetted with MeOH under N₂ atmosphere, 25 mg) under 1 atmosphere H₂ (balloon) for 1.5 hours at room temperature. The catalyst was filtered, the solvent was evaporated in vacuo and the residue was recrystallized from aqueous ethanol. Yield: 127 mg, (0.29 mmol, 81%). R_f = 0.28 (80:20:1 CHCl₃-MeOH-NH₄OH). mp = 136-42° C (becomes translucent). [α]_D = + 7.5° (1.2, MeOH). MS(FAB +) m/z 442(m + H)⁺, 464(m + Na)⁺, 273(-168), 245(-196). ¹H NMR(DMSO_{d6}, 300 MHz) δ 1.83-1.96(m, 1H), 1.98-2.10(m, 1H), 2.30(brt, 2H), 4.50(m, 1H; dd(+ D₂O), J=5.9Hz), 7.05(dt, 1H), 7.12-7.40-(m, 9H), 7.46(d, 1H), 7.63(d, J=7Hz, 1H), 7.68(d, J=5Hz, 1H), 8.55(d, J=7Hz, 1H), 9.30(s, 1H), 11.6(s, 1H), 12.16-(brs, 1H). Analysis calculated for C₂₆H₂₃N₃O₄ H₂O: C 69.32, H 5.37, N 9.33, Found: C 69.58, H 5.27, N, 9.33.

EXAMPLE 39

Boc-(γ -benzyl ester)-R-Glu-(diphenylmethyl)amide

BOPCI (255 mg, 1.0 mmol) was added to a solution of Boc(γ -benzyl ester)-R-Glutamic acid (337 mg, 1.0 mmol) and diphenylmethylamine (431 mL, 2.5 mmol) in dry THF cooled to 4° C. The reaction was allowed to attain room temperature overnight. The reaction mixture was poured into EtOAc and extracted successively with 1 M H₃PO₄ (3x), 5% NaHCO₃ (3x), H₂O (3x) then dried over MgSO₄, filtered and the solvent evaporated in vacuo. The crude product was chromatographed on flash silica gel eluted with 4:1 hexanes/ethyl acetate. Yield: 181 mg, (0.36 mmol), 36%. MS(Cl) m/e 503(m + H)⁺, 520(m + NH₄)⁺, 447(-56), 403(-100), 312(-191). ¹H NMR(CDCl₃, 300MHz) δ 1.42(s, 9H), 1.90-2.03(m, 1H), 2.12-2.23(m, 1H), 2.37-2.6-(m, 2H), 4.21(m, 1H), 5.12(s, 2H), 5.25(brd, J=7Hz, 1H), 6.20(d, J=8Hz, 1H), 7.02(brd, J=7Hz, 1H), 7.20-7.33-(m, 15H). C, H, N Analysis calculated for C₃₀H₃₄N₂O₅: C 71.69, H 6.82, N 5.58; Found: C 71.46, H 6.89, N 5.58.

EXAMPLE 40(γ -Benzyl ester)-R-Glu-(diphenylmethyl)amide hydrochloride

Boc-(γ -benzyl ester)-R-Glu-(diphenylmethyl)amide (100 mg, 0.20 mmol) was dissolved in 4N HCl-dioxane (1.0 mL, 4mmol) under N₂. After 2 hours, the solvent was evaporated. R_f = 0.21 (18:1 CHCl₃-EtOH). Product was used directly in the next step. R_f = 0.80 (90:10:1), CHCl₃-MeOH-NH₄OH).

EXAMPLE 41N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(diphenylmethyl)amide

EDCI (42.1 mg, 0.22 mmol) was added to a cooled (4° C) solution of indole-2-carboxylic acid (36 mg, 0.22 mmol), hydrochloride salt of example 40 (0.2 mmol), HOBT (30 mg, 0.22 mmol) and TEA (62 mL, 0.44 mmol) in 4 mL dry CH₂Cl₂. The reaction was allowed to reach ambient temperature. After 3 days, the reaction mixture was poured into EtOAc and extracted successively with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), H₂O (3x), then dried over MgSO₄, filtered and evaporated. The crude product was recrystallized from EtOAc. Yield: 73 mg, (0.13 mmol), 67%. R_f = 0.19 (2:1 hexanes-ethyl acetate). MS(Cl) m/e 546(m+H)⁺, 455(-91), 438(-108). ¹H NMR(CDCl₃, 300MHz) δ 9.07(s, 1H), 7.66(d, 1H), 7.45(d, 1H), 7.12-7.41(m, 19H), 6.45-(m, 1H), 6.21(dd, 1H, d(+D₂O), J=7Hz), 5.12(d, J=12Hz, 1H), 5.08(d, J=12Hz, 1H), 4.72-4.80(m, 1H), 2.65-2.76-(m, 1H), 2.42-2.52(m, 1H), 2.15-2.32(m, 2H).

EXAMPLE 42N-(2'-Indolylcarbonyl)-R-Glu-N α -(diphenylmethyl)amide

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(diphenylmethyl) amide (60 mg, 0.11 mmol) was dissolved into MeOH (10 mL) and 1 N NaOH (220 μ L) was added at room temperature. After 5 hours, an additional 220 μ L of 1 N NaOH was added and the reaction was refrigerated for 3 days. The solvent was evaporated in vacuo. The residue in H₂O was extracted with EtOAc (3x) then acidified with 6 N HCl and reextracted with EtOAc (3x). The latter EtOAc extractions were combined and washed until neutral to pH paper, then dried over MgSO₄ and evaporated in vacuo. The crude product was recrystallized from aqueous ethanol. R_f = 0.35 (80:20:1 CHCl₃-MeOH-NH₄OH). mp = 224-5° C. [α]_D = 24.4° (c = 0.53, MeOH). MS(FAB+) m/e 456(m+H)⁺, 911(2m+H)⁺. ¹H NMR(CDCl₃, 300MHz) δ 1.95-2.07(m, 2H), 2.24-2.40(m, 2H), 4.62(m, 1H), 6.14-(d, J=8Hz, 1H), 7.03(dt, 1H), 7.18(dt, 1H), 7.22-7.36(m, 11H), 7.43(dd, J=7, \leq 1Hz, 1H), 7.62(d, J=8Hz, 1H), 8.02-(d, J=7Hz, 1H), 8.98(d, J=8Hz, 1H), 11.63(s, 1H), 12.04(brs, 1H). C, H, N Analysis calculated for: C₂₇H₂₅N₃O₄ 0.75 H₂O: C 69.14, H 5.70, N 8.96; Found: C 69.01, H 5.52, N 8.58.

EXAMPLE 43Boc-(γ -benzyl ester)-R-Glu-(2'-indanyl)amide

BOPCl (254 mg, 1.0 mmol) was added to a solution of Boc-(γ -benzyl ester)-R-glutamic acid (337 mg,

1.0 mmol), 2-aminoindane (425 mg, 2.5 mmol) and TEA 348 μ L, 2.5 mmol) in dry THF at room temperature. The reaction mixture was stirred overnight after which the solvent was removed in vacuo. The residue was dissolved into EtOAc and extracted with 1 M H_3PO_4 (3x), 1M Na_2CO_3 (3x), brine (3x) then dried over MgSO_4 , filtered and evaporated in vacuo. The crude product was crystallized from hexanes-
 5 CHCl_3 . Yield: 335 mg, (0.69 mmol, 69%). R_f = 0.26 (2:1 hexanes-ethyl acetate).

EXAMPLE 44(γ -Benzyl ester)-R-Glu-(2'-indanyl)amide hydrochloride

15 Boc-(γ -benzyl ester)-R-Glu-2'-indanylamine (300 mg, 0.61 mmol) was dissolved in HCl-dioxane (3 mL, 12 mmol) at room temperature under N_2 . The solvent was removed after 30 minutes and the crude product was placed under high vacuum overnight.

EXAMPLE 45N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-Glu-(2'-indanyl)amide

EDCI (124 mg, 0.65 mmol) was added to a solution of indole-2-carboxylic acid (105 mg, 0.65 mmol), hydrochloride salt from example 44 (0.61 mmol assumed), HOBt (88 mg, 0.65 mmol) and TEA (181 μ L, 1.3 mmol) in dry CH_2Cl_2 (10 mL). The reaction was stirred overnight then poured into EtOAc and extracted
 30 successively with 1 N H_3PO_4 (3x), 1M Na_2CO_3 (3x), H_2O (2x), brine (1x) then dried over MgSO_4 , filtered and evaporated in vacuo. The crude product was mixed with EtOAc, cooled to -20°C and filtered. Yield: 235 mg, (0.47 mmol, 78%). R_f = 0.17 (2:1 hexanes-EtOAc).

EXAMPLE 46N-(2'-Indolylcarbonyl)-R-Glu-N $^\alpha$ -(2'-indanyl)amide

Benzyl ester of example 45 (215 mg, 0.43 mmol) was dissolved into 2:1 MeOH-EtOAc (30 mL) and treated with 10% palladium on carbon (50 mg, prewetted with MeOH under N_2) under one atmosphere H_2 - (balloon). After 2 hours, the catalyst was filtered and the filtrate was evaporated to dryness in vacuo. The
 45 crude product was recrystallized from 80% aqueous ethanol. R_f 0.21 (80:20:1 CHCl_3 -MeOH- NH_4OH). mp = $184-5^\circ\text{C}$. $[\alpha]_D^{25} = -60.2^\circ$ (c = 1.13, MeOH). MS(FAB+) m/e 406(m+H) $^+$, 289(-117), 273(-133), 245(-161). ^1H NMR($\text{DMSO}-d_6$, 300MHz) δ 1.85-2.05(m, 2H), 2.22-2.39(m, 2H), 2.73-2.85(m, 2H), 3.13-3.23(m, 2H), 4.40-4.52-(m, 2H), 7.04(t, J = 7Hz, 1H), 7.12-7.26(m, 6H), 7.42(d, J = 9Hz, 1H), 7.62(d, J = 8Hz, 1H), 8.38(d, J = 7Hz, 1H), 8.48-(d, J = 8Hz, 1H), 11.59(s, 1H), 12.15(brs, 1H). C,H,N Analysis calculated for: $\text{C}_{23}\text{H}_{23}\text{N}_3\text{O}_4$: C 68.13, H 5.72, N 10.36; Found: C 67.89, H 5.63, N 10.18.

EXAMPLE 47Boc-(γ -benzyl ester)-R-Glu-(3'-quinolyl)amide

BOPCI (254 mg, 1.0 mmol) was added to a cooled (4° C) solution of Boc-(γ -benzyl ester)-R-Glutamic acid (337 mg, 1.0 mmol) and 3-aminoquinoline (361 mg, 2.5 mmol) and TEA (153 μ L, 1.1 mmol), in dry THF (15 mL). The reaction was allowed to reach ambient temperature. After 3 days, additional TEA (153 μ L) and BOPCI (30 mg) was added and stirring continued for 1 day. The solvent was evaporated in vacuo and the residue, dissolved in EtOAc, was extracted successively with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), brine (3x), then dried over MgSO₄, filtered and evaporated. Yield: 344 mg, (0.74 mmol, 74%). R_f = 0.31 (18:1 CHCl₃-EtOH).

EXAMPLE 48

(γ -Benzyl ester)-R-Glu-(3'-quinolyl)amide

Boc-(γ -benzyl ester)-R-Glu-(3'-quinolyl)amide (334 mg, 0.74 mmol) was mixed with HCl-dioxane (3.7 mL, 15 mmol) under an N₂ atmosphere at ambient temperature. Additional HCl-Dioxane (3.7 mL) was added after 1 hour. After 2 hours, the solvent was evaporated and the residue was placed under high vacuum overnight. R_f = 0.10 (18:1 CHCl₃-EtOH).

EXAMPLE 49

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(3'-quinolyl)amide

EDCI (143 mg, 0.75 mmol) was added to a cooled (4° C) solution of indole-2-carboxylic acid (121 mg, 0.75 mmol), hydrochloride salt of example 48 (0.74 mmol assumed), HOBt (101 mg, 0.75 mmol), TEA (209 μ L, 1.5 mmol) in CH₂Cl₂ (15 mL). The reaction was allowed to attain ambient temperature overnight. The solvent was evaporated in vacuo and the residue was dissolved in EtOAc and extracted with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), brine (3x) then dried over MgSO₄, filtered and the solvent evaporated in vacuo. The crude product was recrystallized from EtOAc-hexanes to give: 174 mg, (0.34 mmol) 46%. R_f = 0.22 (1:2 hexanes-EtOAc). mp = 175-7° C. [α]_D = -3.7° (c = 0.83, MeOH). MS(Cl) m/e 507(m+H)⁺, 399(-108), 272(-235), 256(-251). ¹H NMR(CDCl₃, 300 MHz) δ 2.32-2.41(m, 2H), 2.58-2.68(m, 1H), 2.33-2.43(m, 1H), 4.89-(brq, 1H), 5.12(d, J = 14Hz, 1H), 5.17(d, J = 14Hz, 1H), 7.08(d, J < 1Hz, 1H), 7.13(dt, 1H), 7.26(s, 1H), 7.30(s, 5H), 7.41(d, J = 8Hz, 1H), 7.48(dt, J = 1.8Hz, 1H), 7.60(dt, J = 1.7Hz, 1H), 7.68(brt, J = 9Hz, 1H), 7.82(d, J = 7Hz, 1H), 8.02(d, J = 8Hz, 1H), 8.66(d, J = 2Hz, 1H), 8.79(d, J = 2Hz, 1H), 9.37(s, 1H), 9.54(s, 1H). C, H, N Analysis calculated for C₃₀H₂₆N₄O₄ · H₂O: C 69.89, H 5.28, N 10.87; Found: C 69.78, H 5.22, N 10.88.

EXAMPLE 50

N-(2'-Indolylcarbonyl)-R-Glu-N ^{α} -(3'-quinolyl)amide

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(3'-quinolyl)amide (150 mg, 0.30 mmol) was dissolved in MeOH (10 mL) and then treated with 5% palladium on barium sulfate (Pd/BaSO₄) (50 mg) under 1 atmosphere H₂ (balloon). After 4 hours, the catalyst was removed by filtration and the solvent was evaporated in vacuo. The crude product was recrystallized from aqueous methanol to give: 69 mg, (0.16 mmol, 55%). [α]_D = -43.8° (c = 1.04, 1:1 DMF-MeOH). MS(FAB+) m/e 417(m+H)⁺, 307(-110), 277(140), 257(-160). ¹H NMR(DMSO-d₆, 300MHz) δ 2.03-2.25(m, 2H), 2.35-2.5(m, 2H), 4.68(m, 1H; dd(D₂O), J = 5.9Hz), 7.05(dt, J = 1.7Hz, 1H), 7.20(dt, J = 1.7Hz, 1H), 7.32(d, J = 1Hz, 1H), 7.43(d, J = 8Hz, 1H), 7.58(dt, J = 1.8Hz, 1H), 7.63-7.68(m, 2H), 7.95(dt, J = 1.9Hz, 2H), 8.72(d, J = 2Hz, 1H), 8.74(d, J = 8Hz, 1H), 8.97(d, J = 2Hz, 1H), 10.16(s, 1H),

11.12(brs,1H), 12.23(brs,1H). C,H,N Analysis calculated for $C_{23}H_{20}N_4O_4 \cdot H_2O$: C 65.63, H 4.90, N 13.31; Found: C 65.84, H 4.93, N 13.14.

EXAMPLE 51

Boc-(γ -benzyl ester)-R-Glu-(1'-adamantyl)amide

BOPCl (254 mg, 1.0 mmol) was added to a cooled solution (4°C) of Boc-(γ -benzyl ester)-R-Glutamic acid (337 mg, 1.0 mmol), 1-aminoadamantane (378.3 mg, 2.5 mmol), and TEA (153 μL , 1.1 mmol) in dry THF (15 mL). The reaction was allowed to reach ambient temperature overnight. The solvent was evaporated in vacuo and the residue, dissolved in ethyl acetate, was extracted successively with 1 M H_3PO_4 (3x), 1 M Na_2CO_3 (3x), brine (3x) then dried over $MgSO_4$, filtered and evaporated in vacuo. Yield: 276 mg, (0.59 mmol, 59%). R_f = 0.75 (2:1 hexanes-EtOAc). MS(Cl) m/e 471(m+H)⁺, 415(-56), 387(-84), 370(-101). 1H NMR($CDCl_3$, 300MHz) δ 7.33-7.38(m, 5H), 5.78(s, 1H), 5.22(bdd, J = 2.7 Hz, 1H), 5.12(s, 2H), 4.36-4.49(m, 1H), 3.91-4.12(m, 2H), 2.37-2.59(m, 2H), 2.05-2.12(m, 6H), 1.96(bd, J = 2 Hz, 7H), 1.82-1.92(m, 1H), 1.64-1.68(m, 9H).

EXAMPLE 52

(γ -Benzyl ester)-R-Glu-(1'-adamantyl)amide hydrochloride

Boc-(γ -benzyl ester)-R-Glu-(1'-adamantyl)amide (276 mg, 0.59 mmol) was mixed with HCl-dioxane (3.0 mL, 12 mmol) under an N_2 atmosphere at ambient temperature. After 30 minutes, the solvent was evaporated and the residue was placed under high vacuum over night. R_f = 0.70 (80:20:1 $CHCl_3$ -MeOH- NH_4OH).

EXAMPLE 53

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(1'-adamantyl)amide

EDCl (115 mg, 0.6 mmol) was added to a cooled (4°C) solution of indole-2-carboxylic acid (97 mg, 0.60 mmol), hydrochloride salt of example 52 (0.59 mmol assumed), HOBT (81 mg, 0.6 mmol), and TEA (132 μL , 1.2 mmol) in CH_2Cl_2 (10 mL). The reaction was allowed to attain ambient temperature overnight. The solvent was evaporated and the residue was dissolved in EtOAc and extracted with 1 M H_3PO_4 (3x), 1M Na_2CO_3 (3x), brine (3x). The crude product was chromatographed on flash silica gel eluted with a step gradient of 4:1 to 2:1 hexanes-ethyl acetate. Yield: 179 mg, (0.35 mmol), 59%. R_f = 0.24 (2:1 hexanes-ethyl acetate).

EXAMPLE 54

N-(2'-Indolylcarbonyl)-R-Glu- N^α -(1'-adamantyl)amide

N-(2'-Indolylcarbonyl)-(γ-benzylester)-R-Glu-(1'-adamantyl)amide (176 mg, 0.35 mmol) was dissolved in MeOH (10 mL) and then treated with 5% Pd/BaSO₄ (100 mg) under H₂ atmosphere (balloon). After 3 hours, the catalyst was removed by filtration and the solvent was evaporated in vacuo. The crude product was recrystallized from aqueous methanol to give: 117 mg, (0.28 mmol, 79%). [α]_D = -11.6° (c = 0.98, MeOH).
 5 MS(FAB+) m/e 424(m+H)⁺, 408(-16), 273(-151), 246(-178). ¹H NMR(DMSO-d₆, 300 MHz) δ 1.62(brs, 6H), 1.93(brs, 7H), 2.01(brs, 4H), 2.26-2.42(m, 2H), 4.43(m, 1H, + D₂O: dd, J = 6.8 Hz), 7.03(dt, J = 8 Hz, 1), 7.19(dt, J = 1.7 Hz, 1H), 7.23(d, J = 2 Hz, 1H), 7.42(d, J = 8 Hz, 1H), 7.45(s, 1H), 7.52(d, J = 8 Hz, 1H), 8.33(d, J = 8 Hz, 1), 11.11(brs, 1H), 12.17(brs, 1H). C,H,N Analysis calculated for C₂₄H₂₉N₃O₄ · H₂O: C 67.35, H 6.95, N 9.82; Found: C 67.12, H 6.97, N 9.73.

EXAMPLE 55

Boc-(γ-benzyl ester)-R-Glu-(1',2'-diphenylethyl)amide

BOPCl (254 mg, 1.0 mmol) was added to a cooled solution (4° C) of Boc-(γ-benzyl ester)-R-Glutamic acid (337 mg, 1.0 mmol), 1,2-diphenylethylamine (484 μL, 2.5 mmol), and TEA (153 μL, 1.1 mmol) in dry THF (15 mL). The reaction was allowed to reach ambient temperature overnight. After 3 days, the solvent was evaporated in vacuo and the residue, dissolved in ethyl acetate, was extracted successively with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), brine (3x) then dried over MgSO₄, filtered and evaporated in vacuo. The crude product was recrystallized from hexanes-ethyl acetate. Yield: 277 mg, (0.54 mmol, 54%). R_f = 0.46 (2:1 hexanes-ethyl acetate).

EXAMPLE 56

(γ-Benzyl ester)-R-Glu-(1',2'-diphenylethyl)amide hydrochloride

Boc-(γ-benzyl ester)-R-Glu-(1',2'-diphenylethyl)amide (247 mg, 0.48 mmol) was mixed with HCl-dioxane (2.5 mL, 10 mmol) under an N₂ atmosphere at ambient temperature. After 1 hour, the solvent was evaporated and the residue was placed under high vacuum overnight. R_f = 0.63, 0.69 (90:10:1 CHCl₃-MeOH-NH₄OH).

EXAMPLE 57

N-(2'-Indolylcarbonyl)-(γ-benzyl ester)-R-Glu-(1',2'-diphenylethyl)amide

EDCI (95 mg, 0.48 mmol) was added to a cooled (4° C) solution of indole-2-carboxylic acid (77 mg, 0.48 mmol), hydrochloride salt of example 56 (0.48 mmol assumed), HOBt (65 mg, 0.48 mmol), and TEA (139 μL, 1.0 mmol) in anhydrous CH₂Cl₂ (10 mL). The reaction was allowed to attain ambient temperature overnight. After 34 days, the solvent was evaporated and the residue, dissolved in EtOAc, was extracted with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), brine (3x) then dried over MgSO₄, filtered and the solvent evaporated in vacuo. Yield: 253 mg, (0.45 mmol, 94%). R_f = 0.20 (2:1 hexanes-EtOAc). MS(FAB+) m/e 560(m+H)⁺, 363(-197).

EXAMPLE 58

N-(2'-Indolylcarbonyl)-R-Glu-N^α-(1',2'-diphenylethyl)amide

N-(2'-Indolylcarbonyl)-(γ-benzyl ester)-R-Glu-(1',2'-diphenylethyl)amide (176 mg, 0.35 mmol) was dissolved in MeOH (50 mL) with warming and, after returning to ambient temperature, was treated with 1 N NaOH (560 μL, 0.56 mmol). After 3 hours, the solvent was evaporated in vacuo and the residue, dissolved in EtOAc, was extracted with 1 M H₃PO₄ (3x), brine (3x) then dried over MgSO₄, filtered and the solvent evaporated in vacuo. The tlc indicated two slightly resolved spots were present (18:1 CHCl₃-EtOH). The crude product was chromatographed on flash silica gel eluted with 18:1 CHCl₃-EtOH to give 71 mg, (0.15 mmol, 54%). A small amount of each isomer was isolated pure and MS(FAB+) showed them to be methyl esters. The combined product was redissolved in MeOH and again treated with 1 N NaOH (0.5 mL, 0.5 mmol) until both methyl esters were consumed (determined by tlc as above). The solvent was then evaporated in vacuo and the residue, dissolved in H₂O, was acidified with 6 N HCl and extracted into EtOAc (3x). The combined layers were washed with H₂O until neutral to pH paper then dried over MgSO₄, filtered and evaporated in vacuo. The crude product was chromatographed by Prep HPLC (C₁₈ column; with a gradient of 25 - 80% CH₃CN over 10 minutes with 0.05 M NH₄OAc, pH 4.5, as the aqueous buffer). [α]_D = + 7.5° (c = 1.07, DMF). mp = 218-219.5° C. MS(FAB+) m/e 470(m+H)⁺, 284(-186), 273(-197). ¹H NMR-(CD₃OD,300MHz): δ (1:1 mixture of diastereomers) 1.8-2.1(m,2H), 2.14-2.21(m,1H), 2.31(t,J=7Hz,1H), 2.95-3.18(m,2H), 4.06-4.14(m,1H), 5.15(t,J=7Hz,0.5H), 5.22(dd,J=5,10Hz,0.5H), 6.98-7.47(m,14H), 7.08-(d,J=7Hz,0.5H), 7.63(d,J=7Hz,0.5H). C,H,N Analysis calculated for C₂₈H₂₇N₃O₄ 1.5 H₂O: C 67.73, H 6.09, N 8.46; Found: C 67.59, H 5.77, N 8.32.

EXAMPLE 59N-(Nⁱ-Methyl-2'-Indolylcarbonyl)-(γ-benzyl ester)-R-Glu-di-n-pentylamide

EDCI (96 mg, 0.50 mmol) was added to a cooled (4° C) solution of N-methylindole-2-carboxylic acid (88 mg, 0.50 mmol), hydrochloride salt of example 2 (206 mg, 0.50 mmol), HOBt (68 mg, 0.50 mmol), and TEA (140 μL, 1.0 mmol) in CH₂Cl₂ (15 mL). The reaction was allowed to attain ambient temperature overnight. The solvent was evaporated and the residue was dissolved in EtOAc and extracted with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), brine (3x) then dried over MgSO₄, filtered and the solvent evaporated in vacuo. Yield: 262 mg, (0.49 mmol, 98%). R_f = 0.86 (18:1 CHCl₃-EtOH).

EXAMPLE 60N-(Nⁱ-Methyl-2'-Indolylcarbonyl)-R-Glu-N^α,N^α-dipentylamide

The benzyl ester of example 59 (245 mg, 0.46 mmol) was dissolved in MeOH (20 mL) and then treated with 5% Pd/BaSO₄ (50 mg) under 1 atmosphere H₂ (balloon). After 2 hours, the catalyst was removed by filtration and the solvent was evaporated in vacuo. The crude product was dried under high vacuum overnight. R_f = 0.42 (80:20:1 CHCl₃-MeOH-NH₄OH). mp = 132-133° C. [α]_D = + 8.4° (c = 1.09, MeOH). MS(FAB+) m/e 444(m+H)⁺, 287(-157), 269(-175), 259(-185). ¹H NMR(DMSO-d₆,300MHz) δ 0.83-0.87(m,6H), 1.17-1.33(m,8H), 1.41-1.52(m,2H), 1.52-1.70(brm,2H), 1.85-1.94(brm,1H), 2.36(t,J=7Hz,2H), 3.08(quintet,1H), 3.25-3.50(m,3H), 3.96(s,3H), 4.87(m,1H), 7.11(t,J=7Hz,1H), 7.22(s,1H), 7.28(dt,J=1,7Hz,1H), 7.52-(d,J=8Hz,1H), 7.64(d,J=8Hz,1H), 8.54(d,J=8Hz,1H), 12.20(bs,1H). C,H,N Analysis calculated for C₂₅H₃₇N₃O₄: C 67.69, H 8.41, N 9.47; Found: C 67.52, H 8.34, N 9.43.

EXAMPLE 61

Boc-(γ -benzyl ester)-R-Glu-(9'-fluorenyl)amide

BOPCI (254 mg, 1.0 mmol) was added to a cooled solution (4 °C) of Boc-(γ -benzyl ester)-R-Glutamic acid (337 mg, 1.0 mmol), 9-aminofluorene (453 mg, 2.5 mmol), and TEA (153 μ L, 1.1 mmol) in dry THF (15 mL). The reaction was allowed to reach ambient temperature overnight after which additional BOPCI (50 mg) and TEA (30 μ L) were added. After several hours, the solvent was evaporated in vacuo and the residue was dissolved in EtOAc and extracted successively with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), brine (3x) then dried over MgSO₄, filtered and evaporated in vacuo. The crude product was reprecipitated from hexanes-ethyl acetate. Yield: 163 mg, (0.33 mmol, 33%). R_f = 0.41 (2:1 hexanes-EtOAc). MS(Cl) m/e 501(m + H)⁺, 445(-56), 401(-100). ¹H NMR(CDCl₃, 300MHz) δ 7.70(d, J = 7Hz, 2H), 7.52(d, J = 7Hz, 2H), 7.40(t, J = 7Hz, 2H), 7.26-7.32(m, 9H), 6.40(bd, J = 8Hz, 1H), 6.18(d, J = 8Hz, 1H), 5.32(bd, J = 6Hz, 1H), 5.10(s, 2H), 4.18-4.27(m, 1H), 2.45-2.67(m, 2H), 2.18-2.30(m, 1H), 1.98-2.10 (m, 1H), 1.49(s, 9H).

EXAMPLE 62(γ -Benzyl ester)-R-Glu-(9'-fluorenyl)amide hydrochloride

Boc-(γ -benzyl ester)-R-Glu-(9'-fluorenyl)amide (158 mg, 0.32 mmol) was mixed with HCl-dioxane (2.0 mL, 8.0 mmol) under an N₂ atmosphere at ambient temperature. After 90 minutes, the product was precipitated with the addition of ether and the resulting solid collected and dried in vacuo. Yield: 133 mg, (0.31 mmol, 97%). R_f = 0.10 (100:60:18:33 EtOAc-pyridine-acetic acid-H₂O).

EXAMPLE 63N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(9'-fluorenyl)amide

EDCI (69 mg, 0.35 mmol) was added to a cooled (4 °C) solution of indole-2-carboxylic acid (56 mg, 0.35 mmol), hydrochloride salt from example 62 (133 mg, 0.31 mmol), HOBt (47 mg, 0.35 mmol) and TEA (98 μ L, 0.7 mmol) in CH₂Cl₂ (10 mL). The reaction was allowed to attain ambient temperature overnight. The solvent was evaporated and the residue was dissolved in ethyl acetate and extracted with 1 M H₃PO₄ (3x), 1 M Na₂CO₃ (3x), brine (3x). The crude product was recrystallized from ethyl acetate to give: 115 mg, (0.119 mmol), 57%. R_f = 0.62 (1:1 hexanes-ethyl acetate).

EXAMPLE 64N-(2'-Indolylcarbonyl)-R-Glu-N α -(9'-fluorenyl)amide

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(9'-fluorenyl)amide (104 mg, 0.18 mmol) was dissolved in 3:1 MeOH-EtOAc (20 mL) and then treated with 5% Pd/BaSO₄ (50 mg) under 1 atmosphere H₂ (balloon) overnight. The catalyst was removed by filtration and the solvent was evaporated in vacuo. The crude product was recrystallized from aqueous ethanol. R_f = 0.15 (80:20:1 CHCl₃-MeOH-NH₄OH). mp = 265-8 °C. [α]_D = -35.2° (c = 1.07, 1:1 DMF - MeOH). MS(Cl) m/e 454(m + H)⁺, 436(-18), 273(-181). ¹H NMR-(DMSO-d₆, 300MHz) δ 1.98-2.17(m, 2H), 2.37-2.44(m, 2H), 4.52(brq, 1H, + D₂O: dd, J = 5.8Hz), 6.05-6.08(m, 1H), 7.04(t, J = 7Hz, 1H), 7.19(t, J = 7Hz, 1H), 7.27(d, J = 1Hz, 1H), 7.33(d, J = 7Hz, 2H), 7.40-7.52(m, 5H), 7.62(d, J = 8Hz, 1H), 7.36(d, J = 7Hz, 2H), 8.53(d, J = 7Hz, 1H), 8.70(d, J = 8Hz, 1H), 11.63(brs, 1H). C, H, N Analysis

calculated for: $C_{27}H_{23}N_3O_4 \cdot H_2O$: C 70.12, H 5.23, N 9.09; Found: C 70.17, H 5.17, N 9.05.

EXAMPLE 65

Cbz-(γ -benzyl ester)-R-Glu-(1',3'-diphenyl-5'-pyrazyl)amide

Phosphorous pentachloride (PCl_5) (208 mg, 1.0 mmol) was added in one portion to a solution of Cbz-(γ -benzyl ester)-R-Glutamic acid (371 mg, 1.0 mmol) in CH_2Cl_2 (10 mL) at $-20^\circ C$. After 20 minutes, 1,3-diphenyl-5-aminopyrazole (588 mg, 2.5 mmol) was added and the temperature was raised to $0^\circ C$ for 2 hours. The reaction mixture was poured into ethyl acetate and extracted successively with 1 N H_3PO_4 (3x), 1 M Na_2CO_3 (3x), and H_2O (3x) then dried over $MgSO_4$, filtered and the solvent evaporated in vacuo. The residue was chromatographed on flash silica eluted with a stepped gradient from 4:1 to 2:1 hexanes-ethyl acetate. Yield: 555 mg, (0.94 mmol, 74%). R_f = 0.31 (2:1 hexanes-ethyl acetate). 1H NMR($CDCl_3$, 300MHz) δ 8.69(bs, 1H), 7.87(dd, J ~ 1.7 Hz, 2H), 7.30-7.50(m, 18H), 7.05(s, 1H), 5.66(bd, J = 7 Hz, 1H), 5.02-5.11(m, 4H), 4.28-4.34(m, 1H), 2.58-2.68(m, 1H), 2.41-2.52(m, 1H), 2.16-2.26(m, 1H), 1.94-2.06(m, 1H).

EXAMPLE 66

R-Glu- N^α -(1',3'-diphenyl-5'-pyrazyl)amide

Cbz-(γ -benzyl ester)-R-Glu-(1',3'-diphenyl-5'-pyrazyl)amide (417 mg, 0.70 mmol) and 10% Pd/C (100 mg, prewetted with MeOH under an N_2 atmosphere) were combined in ethanol (100 mL) under 1 atmosphere H_2 (balloon) overnight. The mixture was then filtered and rinsed with fresh solvent. Evaporation gave the deprotected product (260 mg, 0.70 mmol, quantitative). R_f = 0.23 (8:1:1 butanol-AcOH- H_2O). The product was used directly in the next step.

EXAMPLE 67

N-(2-Indolylcarbonyl)-R-Glu- N^α -(1',3'-diphenyl-5'-pyrazyl)amide

2-Indolecarboxy-2,4,6-trichlorophenyl ester (255 mg, 0.75 mmol) and R-Glu-(1',3'-diphenyl-5'-pyrazyl)-amide (260 mg, 0.70 mmol) were dissolved in dry ethyl acetate (15 mL) and TEA (209 μ L, 1.5 mmol) was added at room temperature. After one day, additional TEA (209 μ L) was added. After 4 days, the reaction was diluted with ethyl acetate and then extracted successively with 1 M H_3PO_4 (3x), H_2O (3x) and brine (1x) then dried over $MgSO_4$, filtered and evaporated in vacuo. The residue was recrystallized from aqueous ethanol to give: 130 mg, (0.26 mmol, 36%). mp = $261-5^\circ C$ (dec). $[\alpha]_D^{25} = -0.3^\circ$ (c = 1.09, 1:1 DMF-MeOH). MS(FAB+) m/e 508($m+H$) $^+$, 365(-143), 290(-218), 262(-218), 236(-272). 1H NMR($DMSO-d_6$, 300MHz) δ 1.92-2.12(m, 2H), 2.33(brt, J = 7 Hz, 2H), 4.58(m 1H, dd(+ D_2O , J = 5.9 Hz), 6.92(s, 1H), 7.06(dt, J = 1.8 Hz, 1H), 7.21(dt, J = 1.7 Hz, 1H), 7.28(brs, 1H), 7.35(brt, 2H), 7.42-7.47(m, 6H), 7.63(t, J = 8 Hz, 3H), 7.37(d, J = 7 Hz, 2H), 8.78(brd, J = 6 Hz, 1H), 10.28(s, 1H), 11.63(s, 1H). C,H,N Analysis calculated for $C_{29}H_{25}N_5O_4 \cdot H_2O$: C 67.43; H 5.07, N 13.56; Found: C 67.69, H 5.13, N 13.40.

EXAMPLE 68

Boc-(γ -benzyl ester)-R-Glu-(α,α -dimethyl- β -phenylethyl)amide

To a solution of Boc-(γ -benzyl ester)-R-Glutamic acid (337 mg, 1.0 mmol) and α,α -dimethyl- β -phenyl
 5 ethylamine (373 mg, 2.5 mmol) in dry THF (15 mL) was added BOPCl (254 mg, 1.0 mmol) and TEA (1.1
 mmol) and the reaction was stirred for 3 days. After evaporation of the solvent in vacuo, the residue was
 dissolved in ethyl acetate and extracted successively with 1 M H_3PO_4 (3x) 5% NaHCO_3 (3x), brine (3x),
 then dried over MgSO_4 , filtered and the solvent evaporated in vacuo. The oily residue was chromatog-
 10 raphed on flash silica gel eluted with 4:1 hexane-ethyl acetate to yield: 303 mg, (0.65 mmol, 65%) as a
 colorless oil. R_f = 0.46 (2:2 hexanes-ethyl acetate). MS(Cl) m/e 469(m+H)⁺, 413(-56), 369(-100), 278(-191).
 ^1H NMR(CDCl_3 , 300MHz) δ 1.28(s,3H), 1.33(s,3H), 1.42(s,9H), 1.82-1.94(m,1H), 2.03-2.16(m,1H), 2.35-2.54-
 (m,2H), 2.96(d, J=13Hz, 1H), 3.07(d, J=13Hz, 1H), 4.00(m, 1H), 5.12(s, 2H), 5.18(brd, 1H), 5.80(s, 1H), 7.12-
 (dd, J=3,7Hz, 2H), 7.2-7.28(m, 3H), 7.35(s, 5H).

15

EXAMPLE 69

20

(γ -Benzyl ester)-R-Glu-(α,α -dimethyl- β -phenylethyl)amide hydrochloride

The Boc protected material of example 68 (260 mg, 0.56 mmol) was treated with HCl-dioxane (3 mL, 12
 mmol) precooled in an ice bath. An additional 2 mL of HCl-dioxane was added after 2 hours and again at 3
 25 hours at room temperature. After 3.5 hours, the solvent was evaporated and the residue was used directly.
 MS(Cl) m/e 369(m+H)⁺, 351(-18), 277(-92), 236(-133), 192 (-177). R_f = 0.85 (80:20:1 CHCl_3 -MeOH-
 NH_4OH), 0.46 (100:60:18:33 EtOAc-pyridine-HOAc- H_2O).

30

EXAMPLE 70

35

N-(2'-Indolylcarbonyl)-(γ -benzyl ester)-R-Glu-(α,α -dimethyl- β -phenylethyl)amide

EDCI (115 mg, 0.6 mmol) was added to a solution of hydrochloride salt of example 69 (0.56 mmol
 assumed), indole-2-carboxylic acid (97 mg, 0.6 mmol), HOBt (81 mg, 0.6 mmol) and TEA (167 μL , 1.2
 mmol) in CH_2Cl_2 (10 mL) cooled to 4°C in an ice bath. The reaction was allowed to warm to ambient
 40 temperature overnight. The solvent was evaporated and the residue dissolved in ethyl acetate. This solution
 was extracted with portions of 1 M H_3PO_4 , 1 M Na_2CO_3 , H_2O , then dried over MgSO_4 , filtered and the
 solvent evaporated. The residue was chromatographed on flash silica gel and eluted with 2:1 hexane-ethyl
 acetate to give 164 mg, (0.32 mmol, 57%). R_f = 0.27 (2:1 hexanes-ethyl acetate).

45

EXAMPLE 71

50

N-(2'-Indolylcarbonyl)-R-Glu-N $^{\alpha}$ -(α,α -dimethyl- β -phenethyl)amide

The benzyl ester of example 70 (164 mg, 0.32 mmol) was treated with 10% Pd/C (50 mg), prewetted
 with MeOH under an N_2 atm) in 10 mL MeOH under 1 atmosphere H_2 (balloon) for 1 hour. After filtration of
 55 the catalyst, the solvent was evaporated in vacuo. The residue was recrystallized from hot 80% aqueous
 ethanol to give 101 mg, (0.24 mmol 75% yield). mp = 109-114°C. $[\alpha]_D^{25}$ = -38.0° (c=0.70, MeOH). MS-
 (FAB+) m/e 422(m+H)⁺, 273(-149), 245(-177). ^1H NMR($\text{DMSO-}d_6$, 300MHz) δ 1.18(s,3H), 1.26(s,3H), 1.85-
 2.05(m,2H), 2.21-2.40(m,2H), 2.90(d, J=13Hz, 1H), 3.07(d, J=13Hz, 1H), 4.44(m, 1H, dd, + (D2O), J=6,9Hz),

7.05(dt,1H), 7.11-7.22(m,6H), 7.28(brs,1H), 7.41(s,1H), 7.44(d,J=8Hz,1H), 7.6 (d,J=8Hz,1H), 8.44(d,1H), 11.63(brs,1H), 12.18(brs,1H). C,H,N Analysis calculated for $C_{24}H_{27}N_3O_4 \cdot 0.1 H_2O$: C 64.92, H 6.70, N 9.46; Found: C 64.72, H 6.50; N 9.37.

EXAMPLE 72

N-Cbz-(γ -benzyl ester)-R-Glu-diphenylamide

PCl₅ (208 mg, 1.0 mmol) was added in one portion to a solution of Cbz(γ -benzyl ester)-R-Glutamic acid (371.4 mg, 1.0 mmol) in CH₂Cl₂ (10 mL) at -20 °C. After 20 minutes, diphenylamine (423 mg, 2.5 mmol) was added and the temperature was raised to 0 °C for 2 hours. The reaction mixture was poured into ethyl acetate and extracted successively with 1 N HCl(3x), 1 M Na₂CO₃(3x), and H₂O(3x) then dried over MgSO₄, filtered and the solvent evaporated in vacuo. The residue was chromatographed on flash silica eluted with a stepped gradient from 4:1 to 2:1 hexanes-ethyl acetate. Yield: 405 mg, (0.77 mmol, 77%). R_f = 0.31 (2:1 hexanes-ethyl acetate). MS(Cl) m/e 522(m+H)⁺, 540(m+NH₄)⁺. ¹H NMR(CDCl₃,300MHz) δ 1.84-2.12-(m,2H), 2.24-2.47(m,2H), 4.63(m,1H), 5.05(s,2H), 5.08(brs,2H), 5.60(d,J=8Hz,1H), 7.2-7.5(m,20H).

EXAMPLE 73

R-Glu-N α ,N α -diphenylamide

Cbz-(γ -benzyl ester)-R-Glu-diphenyl amide (400 mg, 0.77 mmol) and 10% Pd/C (50 mg, prewetted with MeOH under an N₂ atmosphere) were combined in MeOH (15 mL) under 1 atmosphere H₂ (balloon) for 3 hours. The catalyst was then filtered and rinsed with fresh solvent. Evaporation gave the deprotected product (244 mg, 0.75 mmol, 98%). R_f = 0.13 (80:20:1 CHCl₃-MeOH-NH₄OH). The product was used directly in the next step.

EXAMPLE 74

N-(2'-Indolylcarbonyl)-R-Glu-N α ,N α -diphenylamide

2-Indolecarboxy-2,4,6-trichlorophenyl ester (29 mg, 0.084 mmol) and R-Glu-diphenyl amide (25 mg, 0.084 mmol) were dissolved in dry THF (4 mL) and TEA (22 μ L, 0.16 mmol) was added at room temperature. The tic at 3 hours showed no reaction and an additional 22 μ L TEA was added. After 2 days, the THF was evaporated in vacuo and the residue, dissolved in ethyl acetate, was extracted successively with 1 M H₃PO₄(1x), H₂O(3x) and brine (1x) then dried over MgSO₄, filtered and evaporated in vacuo. The crude product was recrystallized from hot ethyl acetate-hexanes to give 12.1 mg, (0.027 mmol, 33% yield). R_f = 0.14 (80:20:1 CHCl₃-MeOH-NH₄OH). MS(FAB+) m/e 442(m+H)⁺, 273(-168), 245(-196). ¹H NMR-(CDCl₃-CD₃OD,300MHz) δ 2.03-2.20(m,2H), 2.38(dd,J=8,15Hz, 2H), 5.02(dd,J=5,8Hz,1H), 7.1(dJ<1Hz,1H), 7.13(d,J=7Hz,1H), 7.25-7.48(m,12H), 7.66(d,J=8Hz,1H). C,H,N Analysis calculated for C₂₆H₂₃N₃O₄ EtOAc: C 69.96; H 5.44, N 9.07; Found: C 69.72, H 5.57, N 8.90.

EXAMPLE 75

N-(3-(3'-Pyridyl)acryloyl)-(γ-benzyl ester)-R-Glu-N^α,N^α-di-n-pentylamide

Hydrochloride salt (0.23 g, 0.56 mmol) of example 2, EDCI 0.12 g, 0.6 mmol; HOBt (0.075 g, 0.56 mmol) and 3-(3-pyridyl)acrylic acid (0.085 g, 0.56 mmol) were stirred at 0° C under nitrogen in 10 mL of CH₂Cl₂. To this mixture was added NMM (0.12 mL, 1.1 mmol). The reaction mixture was stirred overnight and permitted to reach ambient temperature. The reaction mixture was poured into ethyl acetate and extracted twice with portions of 1.0 N hydrochloric acid, and once with saturated sodium bicarbonate solution. The organic solution was dried over magnesium sulfate and filtered. Evaporation of the volatiles in vacuo provided a residue which was purified by chromatography using ethyl acetate/hexane as was elutant provided 0.22 g (78% yield) of product. MS(Cl) m/e 508(m+H)⁺. ¹H NMR(CDCl₃,300MHz) δ 0.85-0.96-(m,6H), 1.19-1.42(m,8H), 1.48-1.70(m,4H), 1.78-1.95(m,2H), 2.40-2.60(m,2H), 5.13(m,3H), 5.02-(d,J=15Hz,1H), 6.98(d,J=9Hz,1H), 7.35(m,6H), 7.6(d,J=15Hz,1H), 7.80(d,J=9Hz,1H), 8.56-(dd,J=1,6Hz,1H), 8.72(d,J=3Hz,1H).

EXAMPLE 76N-(8'-Quinolylsulfonyl)-(γ-benzyl ester)-R-Glu-di-n-pentyl amide

To a mixture of hydrochloride salt (1.0 mmol) of example 2 and triethylamine (2.0 mmol) stirring in 10 mL of methylene chloride at 0° C was added 8-quinolylsulfonyl chloride (1.0 mmol). The reaction mixture was allowed to warm to ambient temperature and continue stirring overnight. The reaction mixture was poured into ethyl acetate and extracted with 1.0 N hydrochloric acid and saturated sodium bicarbonate solution. The mixture was dried over magnesium sulfate and filtered. Evaporation of the volatiles in vacuo and chromatography of the residue provided product.

EXAMPLE 77N-(8'-Quinolylsulfonyl)-R-Glu-N^α, N^α-di-n-pentyl amide

The product of example 76 was treated with sodium hydroxide solution in methanol as in example 29. Extraction of the acidified solution numerous times with ethyl acetate and subsequent drying, filtration, concentration, and trituration as in example 29 provided product.

EXAMPLE 78Ethyl N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-Glutamyl (N-benzyl)glycinate

N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-glutamic acid (5.17 grams, 15.3 mmol) and N-benzylglycine ethyl ester (9.12 grams, 47.2 mmol) were stirred in 100 mL of CH₂Cl₂ at -10° to 0° C under nitrogen. To this solution was added BOPCI (4.17 grams, 16.4 mmol) and the reaction mixture allowed to slowly warm to room temperature overnight. The mixture was taken up in ethyl acetate and extracted three times with dilute hydrochloric acid, once with saturated sodium bicarbonate solution and dried over magnesium sulfate. The solution was filtered and concentrated in vacuo. The residue was purified by chromatography using ethyl acetate and hexane as elutants to provide 6.59 grams of pure product (84% yield). MS(Cl) m/e 513(m+H)⁺, 322, 305, 276. ¹H NMR(CDCl₃,300MHz) δ 7.25-7.40 (m,8H), 7.18-7.25(m,2H), 5.36(m,2H), 5.10(m,2H), 4.95-

(bd, J = 15Hz, 1H), 4.53(bd, J = 15Hz, 1H), 4.15(m, 2.5H), 3.70(d, J = 17Hz, 0.5H), 2.40-2.60(m, 2H), 2.20(m, 1H), 1.82(m, 1H), 1.43(bs, 9H), 1.23(m, 3H).

EXAMPLE 79

Ethyl (γ -benzyl ester)-R-Glutamyl(N-benzyl)glycinate hydrochloride

The ethyl ester product of example 78 (3.46 grams, 6.76 mmol) was stirred at 5° C in 10 mL of 4.5 N hydrochloric acid in dioxane. When the reaction was complete by tlc, diethyl ether was added and the solvents removed in vacuo. The product was collected via filtration using diethyl ether. Yield 2.96 grams (97%). MS(FAB) m/e 413(m + H)⁺, 395, 378, 323, 305, 289. ¹H NMR(DMSO-d₆, 300MHz) δ 8.38(bs, 1H), 7.25-7.40(m, 10H), 5.11(bs, 2H), 4.97(bd, J = 17Hz, 0.67H), 4.70(bd, J = 17Hz, 0.33H), 4.35-4.55(m, 3H), 4.22(m, 1H), 4.08(m, 2H), 3.69(d, J = 17Hz, 1H), 3.35(bs, H₂O), 2.60-2.70(m, 2H), 2.10(m, 2H), 1.15(m, 3H).

EXAMPLE 80

Ethyl N-(2'-Naphthoyl)-(γ -benzyl ester)-R-Glutamyl(N-benzyl)glycinate

The hydrochloride of example 79 (288 mg, 0.64 mmol) was coupled to 2-naphthoic acid in a similar manner to that in example 70 to provide product (322 mg, 88% yield) after purification by chromatography. mp = 103-105° C. [α]_D = -6.8° (c = 0.25, MeOH-DMF 1:1). MS(Cl) m/e 567(m + H)⁺, 374, 346. ¹H NMR-(CDCl₃, 300MHz) δ 8.35(s, 1H), 7.85-7.95(m, 4H), 7.56(m, 2H), 7.20-7.40(m, 10H), 5.49(dt, J = 3.5, 8.5Hz, 0.67H), 5.20(m, 0.33H), 5.11(m, 2H), 5.03(d, J = 16Hz, 0.67H), 4.69(s, 0.67H), 4.62(d, J = 16Hz, 0.67), 4.10-4.45(m, 4H), 3.74(d, J = 16Hz, 1H), 2.30-2.70(m, 3H), 2.00-2.10(m, 1H), 1.23(m, 3H). C, H, N analysis calculated for C₃₄H₃₄N₂O₆: C 72.07, H 6.05, N 4.94; found C 71.83, H 6.11, N 4.87.

EXAMPLE 81

Ethyl N-(2-Naphthoyl)-R-Glutamyl(N-benzyl)glycinate

The product of example 80 (167 mg) was subjected to hydrogenolysis as in example 21 utilizing cyclohexadiene as the reductant. Product was purified by chromatography over silica with 0.5% acetic acid in ethyl acetate solution and recovered by lyophilization to provide 121 mg (86% yield) of product. mp = 56-62° C. [α]_D = -10.5° (c = 0.76, MeOH). MS(Cl) m/e 477(m + H)⁺, 284, 194, 155. ¹H NMR-(DMSO-d₆, 300MHz) δ 8.92(bd, J = 7.5Hz, 0.33H), 8.83(bd, J = 7.5Hz, 0.67H), 8.52(bd, J = 5.5Hz, 1H), 8.00(m, 4H), 7.63(m, 2H), 7.46(bd, J = 7Hz, 1H), 7.23-7.40(m, 4H), 5.12(m, 1H), 5.07(bs, 0.33H), 4.95(m, 0.67H), 4.55(m, 2H), 4.19(m, 0.67H), 4.06(q, J = 7Hz, 1.33H), 3.96(m, 0.33H), 3.64(bd, J = 17Hz, 0.67H), 2.43(m, 1.33H), 2.35(m, 0.67H), 1.90-2.10(m, 2H), 1.16(t, J = 7Hz, 2H), 1.09(t, J = 7Hz, 1H). C, H, N analysis calculated for C₂₇H₂₈N₂O₆ 0.25 H₂O: C 67.42, H 5.97, N 5.82; found C 67.43, H 5.88, N 5.77.

EXAMPLE 82

Ethyl N-(3'-Quinolylcarbonyl)-(γ -benzylester)-R-Glutamyl(N-benzyl)glycinate

The hydrochloride of example 79 (325 mg, 0.67 mmol) was coupled to quinoline-3-carboxylic acid in a similar manner to that in example 70 to provide product (331 mg, 87.7% yield) after purification by chromatography. $[\alpha]_D^{25} = +5.5^\circ$ ($c=0.86$, MeOH). MS(Cl) m/e 568($m+H$)⁺, 375, 342. ¹H NMR-(CDCl₃, 300MHz) δ 9.34(d, $J=2$ Hz, 1H), 8.60(d, $J=1.8$ Hz, 1H), 8.17(d, $J=7.5$ Hz, 1H), 7.92(m, 1H), 7.82(m, 1H), 7.63(m, 1H), 7.50(m, 1H), 7.20-7.40(m, 9H), 5.50(dt, $J=1.4, 5$ Hz, 0.67H), 5.00-5.12(m, 3H), 4.69(s, 0.67H), 4.62-(d, $J=16$ Hz, 0.67H), 4.10-4.45(m, 3H), 3.74(d, $J=16$ Hz, 1H), 2.50-2.70(m, 2H), 2.30-2.50(m, 1H), 2.03-2.11(m, 1H), 1.60(bs, H₂O), 1.26(m, 3H). C, H, N analysis calculated for C₃₃H₃₃N₃O₆ 0.7 H₂O: C 8.31, H 5.98, N 7.24; found C 68.28, H 5.88, N 7.13.

EXAMPLE 83

Ethyl N-(3'-Quinolylcarbonyl)-R-Glutamyl(N-benzyl)glycinate

The product of example 82 (100 mg) was subjected to hydrogenolysis as in example 21 utilizing cyclohexadiene as the reductant. Product was purified by chromatography over silica with 0.5% acetic acid in ethyl acetate solution and recovered by lyophilization to provide 55.3 mg (66% yield) of product. mp = 70-79° C. $[\alpha]_D^{25} = -30.4^\circ$ ($c=0.23$, acetone). MS(Cl) m/e 478($m+H$)⁺, 287, 257, 194, 156. ¹H NMR-(DMSO-d₆, 300MHz) δ 9.29(d, $J=2$ Hz, 0.33), 9.25(d, $J=2$ Hz, 0.67H), 9.15(bd, $J=7.5$ Hz, 0.33), 9.05-(bd, $J=7.5$ Hz, 0.67H), 8.91(d, $J=2$ Hz, 0.33H), 8.85(d, $J=2$ Hz, 0.67H), 8.10(m, 2H), 7.88(m, 1H), 7.72(m, 1H), 7.45-(bd, $J=7$ Hz, 1H), 7.22-7.37(m, 4H), 5.13(m, 1.33H), 4.96(m, 0.67H), 4.58(m, 2H), 4.22(m, 0.67H), 4.08-(q, $J=7$ Hz, 1.33H), 3.96(m, 0.33H), 3.65(bd, $J=17$ Hz, 0.67H), 3.35(bs, H₂O), 2.33-2.48(m, 2H), 1.92-2.12(m, 2H), 1.17(t, $J=7$ Hz, 2H), 1.09(t, $J=7$ Hz, 1H). C, H, N analysis calculated for C₂₆H₂₇N₃O₆ 0.33 H₂O: C 64.59, H 5.77, N 8.69; found C 64.47, H 5.69, N 8.71.

EXAMPLE 84

Ethyl N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-(γ-benzylester)-R-Glutamyl(N-benzyl)glycinate

The hydrochloride of example 79 (288 mg, 0.64 mmol) was coupled to 4,8-dihydroxyquinoline-2-carboxylic acid in a similar manner to that in example 20 to provide product (234 mg, 60% yield) after purification by chromatography. mp = 71-74° C. $[\alpha]_D^{25} = -10.7^\circ$ ($c=0.56$, acetone). MS(Cl) m/e 600($m+H$)⁺, 407, 379. ¹H NMR(CDCl₃, 300MHz) δ 7.76(m, 1H), 7.55(m, 1H), 7.20-7.45(m, 11H), 6.95(m, 1H), 5.25(m, 1H), 5.18(bs, 2H), 5.07(bd, $J=16$ Hz, 1H), 4.70-4.90(m, 1H), 4.62(dd, $J=4.5, 16$ Hz, 1H), 4.53(bd, $J=16$ Hz, 1H), 4.18-(m, 2H), 3.69(bd, $J=16$ Hz, 1H), 2.55-2.85(m, 2H), 2.50-2.68(m, 1H), 2.44(m, 2H), 1.23(bt, $J=7.5$ Hz, 3H).

EXAMPLE 85

Ethyl N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-R-Glutamyl(N-benzyl)glycinate

The product of example 84 (57 mg, 0.095 mmol) was subjected to hydrogenolysis as in example 21 utilizing cyclohexadiene as the reductant and DMF (1 mL) to aid in solubilization of the substrate. Product was purified by chromatography over silica with 0.5% acetic acid in ethyl acetate solution and recovered by lyophilization to provide 30.3 mg (60% yield) of product. mp = 103-109° C. $[\alpha]_D^{25} = -10.0^\circ$ ($c=0.13$, MeOH). $[\alpha]_D^{25} = -10.0^\circ$ ($c=0.13$, MeOH). MS(Cl) m/e 510($m+H$)⁺, 483, 464, 305. ¹H NMR(DMSO-d₆, 300MHz)

δ 10.12(bs,1H), 9.72(m,1H), 7.96(bd,J=7Hz,0.67H), 7.69(bd,J=7.5Hz,0.33H), 7.54(m,1H), 7.37-7.50(m,3H), 7.19-7.33(m,4H), 7.12(bd,J=7.5Hz,1H), 5.18(m,0.33H), 4.95(m,0.67H), 4.55(bd,J=7.5Hz,1H), 4.50(m,1H), 4.19(m,0.67H), 4.07(bq,J=7Hz,1.33H), 3.90(m,0.33H), 3.70(bd,J=16Hz,0.67H), 3.34bs,H₂O), 2.25-2.37-(m,2H), 2.15(m,1H), 2.00(m,1H), 1.14(t,J=7Hz,2H), 1.07(t,J=7Hz,1H).

EXAMPLE 86

N-(2'-Methylphenylaminocarbonyl)-(γ-benzyl ester)-S-Glu-di-n-pentylamide

To a solution of the hydrochloride salt of (γ-benzyl ester)-S-Glu-di-n-pentylamide (0.1 g, 0.24 mmol) in THF (7 mL) was added TEA (40 μL, 0.3 mmol) and a slight excess of 1-methylphenylisocyanate (0.55 g, 0.42 mmol). The reaction mixture was stirred overnight at ambient temperature. The solvents were evaporated and residue purified by chromatography using EtOAc and hexane as the elutant mixture. Pure fractions were pooled and the volatiles removed in vacuo to yield an oily product (0.075 g) in 62% yield. MS(Cl) m/e 510(m+H)⁺. ¹H NMR(CDCl₃,300MHz) δ 0.8-0.95(m,6H), 1.1-1.35(m,8H), 1.36-1.5(m,4H), 1.72-1.83(m,1H), 2.0-2.12(m,1H), 2.21(s,3H), 2.38-2.60(m,2H), 2.97-3.08(m,1H), 3.15-3.3(m,1H), 3.4-3.55(m,2H), 4.9(m,1H), 5.1(s,2H), 6.02(d,J=9Hz,1H), 6.47(s,1H), 7.05(t,J=7Hz,1H), 7.08-7.19(m,2H), 7.35(m,5H), 7.48-(d,J=9Hz,1H). C,H,N analysis calculated for C₃₀H₄₃N₃O₄ 0.25 H₂O: C 70.07, H 8.53, N 8.17; found C 69.79, H 8.30, N 8.18.

EXAMPLE 87

Methyl (γ-benzyl ester)-R-Glutamyl-S-phenylglycinate hydrochloride

The product of example 35 was deprotected in a similar manner to that in example 79. The product was concentrated under vacuum to an oil which was used without further purification.

EXAMPLE 88

Methyl N-(2'-Naphthoyl)-(γ-benzyl ester)-R-Glutamyl-S-phenylglycinate

The hydrochloride of example 87 (157 mg, 0.37 mmol) was coupled with 2-naphthoic acid as in example 80 to provide product (156.2 mg, 78% yield) after chromatography. mp = 154-6°C. [α]_D = +38.4° (c=0.24, acetone). MS(Cl) m/e 539(m+H)⁺, 374, 347, 155. ¹H NMR(CDCl₃,300MHz) δ 8.36(s,1H), 7.89(m,4H), 7.56(m,3H), 7.43(d,J=7Hz,1H), 7.32-7.40(m,10H), 5.55(d,J=7Hz,1H), 5.12(s,2H), 4.84-(dt,J=4.8,7.5Hz,1H), 3.70(s,3H), 2.72(m,1H), 2.45(m,1H), 2.10-2.33(m,2H). C,H,N analysis calculated for C₃₂H₃₀N₂O₆: C 69.05, H 5.79, N 5.03; found C 69.26, H 5.74, N 4.90.

EXAMPLE 89

Methyl N-(2'-Naphthoyl)-R-Glutamyl-S-phenylglycinate

The product of example 88 (73 mg, 0.136 mmol) was subjected to hydrogenolysis as in example 21 using cyclohexadiene as the reductant and DMF (1 mL) to solubilize the substrate for transfer to the reaction vessel. Yield 48 mg (79%). mp = 56.63° C. $[\alpha]_D^{25} = +32.9^\circ$ (c=0.34, MeOH). MS(Cl) m/e 449-(m+H)⁺, 329, 284, 273, 257, 155. ¹H NMR(DMSO-d₆, 300MHz) δ 8.90(d, J=7Hz, 1H), 8.64(d, J=8Hz, 1H), 8.52-(bs, 1H), 7.95-8.06(m, 4H), 7.58-7.67(m, 2H), 7.35-7.42(m, 6H), 5.48(d, J=7Hz, 1H), 4.68(m, 1H), 3.63(s, 3H), 2.32-(m, 2H), 1.98(m, 2H).

EXAMPLE 90

Methyl N-(3'-Quinolylcarbonyl)-(γ-benzyl ester)-R-Glutamyl-S-phenylglycinate

The hydrochloride of example 87 (221 mg, 0.528 mmol) was coupled with quinoline-3-carboxylic acid as in example 80 to provide product (170 mg, 63% yield) after chromatography. mp = 136-137.5° C. $[\alpha]_D^{25} = +37.9^\circ$ (c=0.24, acetone). MS(Cl) m/e 540(m+H)⁺, 480, 432, 375, 156. ¹H NMR(CDCl₃, 300MHz) δ 9.34(d, J=2Hz, 1H), 8.62(d, J=2Hz, 1H), 8.18(d, J=7.5Hz, 1H), 7.90(dd, J=1, 7.5Hz, 1H), 7.83(m, 1H), 7.70-(bd, J=7Hz, 1H), 7.6(m, 1H), 7.54(bd, J=7Hz, 1H), 7.28-7.40(m, 10H), 5.55(d, J=7Hz, 1H), 5.16(d, J=12Hz, 1H), 5.12(d, J=12Hz, 1H), 4.85(dt, J=5, 7Hz, 1H), 3.71(s, 3H), 2.75(m, 1H), 2.48(m, 1H), 2.15-2.33(m, 2H). C, H, N analysis calculated for C₃₁H₂₉N₃O₆: C 69.01, H 5.42, N 7.79; found C 68.63, H 5.46, N 7.76.

EXAMPLE 91

Methyl N-(3'-Quinolylcarbonyl)-R-Glutamyl-S-phenylglycinate

The product of example 90 (67 mg, 0.124 mmol) was subjected to hydrogenolysis as in example 21 using cyclohexadiene as the reductant and DMF (1 mL) to solubilize the substrate for transfer to the reaction vessel. Yield 28 mg (50%). mp = 182-185° C. $[\alpha]_D^{25} = +46.6^\circ$ (c=0.075, MeOH). MS(Cl) m/e 450-(m+H)⁺, 285, 256, 156. ¹H NMR(DMSO-d₆, 300MHz) δ 9.30(d, J=2Hz, 1H), 8.96(d, J=7Hz, 1H), 8.90(m, 2H), 8.11(m, 2H), 7.88(m, 1H), 7.71(m, 1H), 7.39(m, 5H), 5.49(d, J=7Hz, 1H), 4.72(m, 1H), 3.64(s, 3H), 3.35(bs, H₂O), 2.32(m, 2H), 1.98(m, 2H). C, H, N analysis calculated for C₂₄H₂₃N₃O₆ 0.67 H₂O: C 62.48, H 5.31, N 9.11; found C 62.23, H 5.29, N 8.93.

EXAMPLE 92

Methyl N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-(γ-benzyl ester)-R-Glutamyl-S-phenylglycinate

The hydrochloride of example 87 (146 mg, 0.35 mmol) was coupled with 4,8-dihydroxyquinoline-2-carboxylic acid as in example 80 to provide product 65.6 mg, (33% yield) after chromatography. mp = 234-5° C. MS(Cl) m/e 572(m+H)⁺, 385, 294, 277, 211, 192, 166. ¹H NMR(DMSO-d₆, 300MHz) δ 10.08(bs, 1H), 9.56(d, J=7Hz, 1H), 9.00(d, J=7Hz, 1H), 7.55(m, 2H), 7.25-7.45(m, 10H), 7.12(d, J=7Hz, 1H), 5.48(d, J=7Hz, 1H), 5.05(s, 2H), 4.75(m, 1H), 3.61(s, 3H), 3.30(bs, H₂O), 2.43(m, 2H), 2.10(m, 2H). C, H, N analysis calculated for C₃₁H₂₉N₃O₈ 0.5 H₂O: C 64.13, H 5.21, N 7.24; found C 64.43, H 5.07, N 7.25.

EXAMPLE 93

Methyl N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-R-Glutamyl-S-phenylglycinate

The product of example 92 (49.7 mg, 0.087 mmol) was subjected to hydrogenolysis as in example 21 using cyclohexadiene as the reductant and DMF (2 mL) to solubilize the substrate for transfer to the reaction vessel. Yield 31.4 mg (75%). MS(Cl) m/e 482(m+H)⁺, 464, 317, 289, 277, 205. ¹H NMR-(DMSO-d₆,300MHz) δ 9.56(m,1H), 8.96(bd,J=7Hz,1H), 7.57(bd,J=8Hz,1H), 7.50(bs,1H), 7.33-7.45(m,7H), 7.12(bd,J=8Hz,1H), 5.47(d,J=7Hz,1H), 4.71(m,1H), 3.62(s,3H), 3.3(bs,H₂O), 2.28(m,2H), 2.00(m,2H). C,H,N analysis calculated for C₂₄H₂₃N₃O₈ 0.33 H₂O: C 59.14, H 4.89, N 8.62; found C 59.07, H 4.95, N 8.39.

EXAMPLE 94Methyl N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-Glutamyl-S-leucinate

N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-glutamic acid (1.077 grams, 3.0 mmol) and methyl-S-leucinate hydrochloride were coupled in a manner similar to that in example 21 to provide 1.28 grams (86% yield) of product after chromatography. mp = 69-71 °C. [α]_D = -1.1° (c=0.62, acetone). MS(Cl) m/e 482(m+NH₄)⁺, 465(m+H)⁺, 409, 365, 309, 292, 274, 257. ¹H NMR(CDCl₃,300MHz) δ 7.36(m,5H), 6.55(m,1H), 5.30(m,1H), 5.14(s,2H), 4.56(m,1H), 4.19(m,1H), 3.72(s,3H), 2.56(dt,J=7,17Hz,1H), 2.47(dt,J=7,17Hz,1H), 2.17(m,1H), 1.97(m,1H), 1.50-1.67(m,3H), 1.41(s,9H), 0.92(d,J=6Hz,6H). C,H,N analysis calculated for C₂₄H₃₅N₂O₇: C 62.05, H 7.81, N 6.03; found C 61.98, H 7.80, N 5.96.

EXAMPLE 95Methyl (γ-benzyl ester)-R-Glutamyl-S-leucinate hydrochloride

The product of example 94 (1.07 grams, 2.30 mmol) was treated with 6.0 mL of 4.5 N hydrochloric acid in dioxane. The hydrochloride was isolated as an oil as in example 87, yield 914 mg (99%). MS(Cl) m/e 365(m+H)⁺, 192. ¹H NMR(DMSO-d₆,300MHz) δ 9.05(d,J=7Hz,1H), 8.34(bs,3H), 7.32-7.42(m,5H), 5.11(s,2H), 4.28-4.36(m,1H), 3.91(t,J=6Hz,1H), 3.64(s,3H), 2.36-2.58(m,2H), 2.02-2.09(m,2H), 1.49-1.65(m,3H), 0.86-(d,J=6Hz,3H), 0.82(d,J=6Hz,3H).

EXAMPLE 96Methyl N-(2'-naphthoyl)-(γ-benzyl ester)-R-Glutamyl-S-leucinate

Naphthalene-2-carboxylic acid (121 mg, 0.70 mmol) and the product of example 95 (200 mg, 0.50 mmol) were coupled and extracted as in example 80. The resulting residue was chromatographed on silica gel eluted with a step gradient of 9:1 to 2:1 hexanes-EtOAc to give 149 mg, 0.29 mmol (44% yield). mp = 95-98 °C. [α]_D = +17.3° (c=0.55, CHCl₃). MS(Cl) m/e 519(m+H)⁺, 428, 411. ¹H NMR(CDCl₃,300MHz) δ 0.94(dd,J=2,6Hz,6H), 1.55-1.71(m,2H), 2.13-2.38(m,3H), 2.55(dt,J=7,17Hz,1H), 2.72-2.82(m,1H), 3.67(s,3H), 4.54-4.65(m,1H), 4.78-4.84(m,1H), 5.13(s,2H), 6.91(d,J=8Hz, 1H), 7.28-7.33(m,5H), 7.50-7.61(m,3H), 7.86-7.94(m,4H), 8.36(s,1H). C,H,N analysis calculated for C₃₀H₃₄N₂O₆: C 69.48, H 6.61, N 5.40; found C 69.21, H 6.71, N 5.33.

EXAMPLE 97

Methyl N-(2'-Naphthoyl)-R-Glutamyl-S-leucinate

5

The product of example 96 (74 mg, 0.14 mmol) was dissolved in 10 mL MeOH and treated with 20 mg of 10% Pd/C (prewetted with solvent under N₂) and 1 atmosphere of hydrogen gas. After 3 hours, the catalyst was filtered and rinsed with fresh solvent. The residue was dissolved in MeOH, filtered through a 0.5 μ filter and then diluted with water. The aqueous mixture was frozen and lyophilized to give 31 mg, 0.072 mmol (52% yield). R_f = 0.22 (80:20:1 CHCl₃-MeOH-NH₄OH). mp = 58-65° C. [α]_D = -31.8° (c=1.0, MeOH). MS(Cl) m/e 429(m+H)⁺, 284. ¹H NMR(DMSO-d₆, 300MHz) δ 0.83(d, J=6Hz, 3H), 0.86(d, J=6Hz, 3H), 1.43-1.63(m, 3H), 1.94-2.03(m, 2H), 2.33(bt, J=7Hz, 2H), 3.62(s, 3H), 4.28-4.36(m, 1H), 4.56(q, J=7Hz, 1H), 7.56-7.65(m, 2H), 7.96-8.06(m, 4H), 8.42(d, J=8Hz, 1H), 8.52(s, 1H), 8.83(d, 7Hz, 1H). C,H,N analysis calculated for C₂₃H₂₈N₂O₆ 0.6 H₂O: C 62.89, H 6.70, N 6.38; found C 62.75, H 6.51, N 6.44.

10

15

EXAMPLE 98

Methyl N-(3'-Quinolylcarbonyl)-(γ -benzyl ester)-R-Glutamyl-S-leucinate

20

25

Quinoline-3-carboxylic acid (139 mg, 0.8 mmol) and the product of example 95 (200 mg, 0.50 mmol) were coupled as in example 82. Silica gel chromatography of the residue eluted with a step gradient of 9:1 to 1:1 hexane-EtOAc gave 99 mg, 0.19 mmol (39% yield). R_f = 0.22 (1:1 hexane-EtOAc). mp = 105-107° C. [α]_D = +16.9° (c=1.5, CHCl₃). MS(Cl) m/e 520(m+H)⁺, 429, 412. ¹H NMR(CDCl₃, 300MHz) δ 0.93-(d, J=2Hz, 3H), 0.96(d, J=2Hz, 3H), 1.55-1.71(m, 3H), 2.17-2.37(m, 2H), 2.57(ddd, J=5.7, 17Hz, 1H), 2.80-(ddd, J=5.8, 17Hz, 1H), 3.69(s, 3H), 4.58-4.65(m, 1H), 4.76-4.82(m, 1H), 5.13(d, J=12Hz, 1H), 5.18-(d, J=12Hz, 1H), 6.90(d, J=8Hz, 1H), 7.28-7.35(m, 5H), 7.63(dt, J=1.8Hz, 1H), 7.76(d, J=7Hz, 1H), 7.82-(dt, J=1.7Hz, 1H), 7.91(d, J=8Hz, 1H), 8.16(d, J=8Hz, 1H), 8.62(d, J=2Hz, 1H), 9.36(d, J=2Hz, 1H). C,H,N analysis calculated for C₂₉H₃₃N₃O₆ 0.3 H₂O: C 66.35, H 6.45, N 8.00; found C 66.34, H 6.43, N 7.96.

30

35

EXAMPLE 99

Methyl N-(3'-Quinolylcarbonyl)-R-Glutamyl-S-leucinate

40

The product of example 98 (100 mg, 0.192 mmol) was treated with 25 mg of 10% Pd/C (prewetted with EtOH under N₂) and 1 atmosphere of hydrogen gas in 25 mL of EtOH. Another 20 mg of catalyst was carefully added after 4 hours and the reaction was allowed to continue overnight. After filtration of the catalyst and evaporation of the solvent, the residue was chromatographed on silica gel eluted with a 9:1 CH₂Cl₂-EtOH to 80:20:1 CHCl₃-MeOH-NH₄OH step gradient. After evaporation of the product fractions, the residue was mixed with aqueous EtOH, frozen and lyophilized to give 39 mg, 0.092 mmol (48% yield). mp = 194-6° C. [α]_D = -26.6° (c=1.64, MeOH). MS(Cl) m/e 430(m+H)⁺, 412, 285, 257, 199, 174, 156. ¹H NMR(CDCl₃, 300 MHz) δ 0.95(dd, J=5.6Hz, 6H), 1.62-1.72(m, 3H), 2.08-2.27(m, 2H), 2.49-2.53(m, 2H), 3.70-(s, 3H), 3.98-4.02(m, 1H), 4.22(dd, J=6.8Hz, 1H), 7.21(dt, J=1.8Hz, 1H), 7.39(dt, J=1.7Hz, 1H), 8.06-8.12(m, 2H), 8.86(d, J=2Hz, 1H), 9.28(d, J=2Hz, 1H).

45

50

55

EXAMPLE 100

Methyl N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-(γ-benzyl ester)-R-Glutamyl-S-leucinate

4,8-Dihydroxyquinoline-2-carboxylic acid (62 mg, 0.3 mmol), the product of example 95 (100 mg, 0.25 mmol), HOBt (41 mg, 0.3 mmol), and TEA (42 μL, 0.3 mmol) were dissolved in 5 mL DMF and treated with EDCI (57 mg, 0.3 mmol). After 3 days, the solvent was evaporated and the residue in EtOAc was extracted as in example 82. The residue from evaporation was chromatographed in silica gel eluted with a 1% to 5% EtOH in CH₂Cl₂ to give 108 mg, 0.196 mmol (78% yield) as a light yellow solid. mp = 208-10 °C. $[\alpha]_D^{25} = -28.0^\circ$ (c=0.46, MeOH). MS(FAB+) m/e 552(m+H)⁺, 374. ¹H NMR(CDCl₃-CD₃OD, 300MHz) δ 0.92-(bd, J=5Hz, 6H), 1.55-1.68(m, 3H), 2.15-2.37(m, 2H), 2.48-2.66(m, 2H), 3.70(s, 3H), 4.52-4.57(m, 1H), 4.73-(dd, J=5, 12Hz, 1H), 5.13(s, 2H), 7.16(bd, J=7Hz, 1H), 7.28-7.33(m, 6H), 7.40(s, 1H), 7.72(dd, J=1, 8Hz, 1H). C, H, N analysis calculated for C₂₉H₃₃N₃O₈ 0.25 H₂O: C 62.41, H 5.91, N 7.39.

EXAMPLE 101Methyl N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-R-Glutamyl-S-leucinate

The product of example 100 (50 mg, 0.091 mmol) was treated with 10 mg of 10% Pd/C in 5 mL MeOH as in example 97. After 2 hours, the catalyst was filtered and the solvent was evaporated. mp = 228-30 °C. $[\alpha]_D^{25} = -42.0^\circ$ (c=1.07, DMF). MS(Cl) m/e 462(m+H)⁺. ¹H NMR(DMSO-d₆, 300MHz) δ 0.83(d, J=6Hz, 3H), 0.88(d, J=6Hz, 3H), 1.43-1.66(m, 3H), 1.42-2.15(m, 2H), 2.32(bt, J=7Hz, 2H), 3.61(s, 3H), 4.29-4.36(m, 1H), 4.56-4.63(m, 1H), 7.10(dd, J=1, 7Hz, 1H), 7.40(t, J=8Hz, 1H), 7.47(bs, 1H), 7.56(dd, J=1, 8Hz, 1H), 8.45(d, J=7Hz, 1H), 9.55(bd, J=7Hz, 1H). C, H, N analysis calculated for C₂₂H₂₇N₃O₈ 1.2 H₂O: C 54.70, H 6.13, N 8.70; found C 54.63, H 5.79, N 8.30.

EXAMPLE 102N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-Glutamyl-bis(carboethoxymethyl)amide

N-(t-Butyloxycarbonyl)-(γ-benzyl ester)-R-glutamic acid (1.98 gram, 5.9 mmol) and diethyl imidodiacetate (5.32 gram, 28.15 mol) were coupled in a manner similar to that in example 78 to provide 2.62 grams (88% yield) of product after chromatography. $[\alpha]_D^{25} = +16.5^\circ$ (c=1.16, MeOH). MS(Cl) m/e 526-(m+NH₄)⁺, 509(m+H)⁺, 470, 453, 436, 423, 409, 380. ¹H NMR(CDCl₃, 300MHz) δ 7.35(m, 5H), 5.31-(bd, J=8Hz, 1H), 5.17(d, J=16Hz, 1H), 5.12(d, J=16Hz, 1H), 4.70(dt, J=4, 9Hz, 1H), 4.38(bs, 2H), 4.05-4.27-(m, 6H), 2.40-2.60(m, 2H), 2.13(m, 1H), 1.73(m, 1H), 1.43(bs, 9H), 1.28(t, J=7Hz, 3H), 1.26(t, J=7Hz, 3H).

EXAMPLE 103(γ-Benzyl ester)-R-Glutamyl-bis(carboethoxymethyl)amide hydrochloride

The product of example 102 (2.16 gram, 4.25 mmol) was deprotected as in example 95 to provide 1.88 grams as an oil which was utilized as is. MS(Cl) m/e 409(m+H)⁺, 391, 380, 363. ¹H NMR(DMSO-d₆, 300MHz) δ 8.46(bs, 3H), 7.30-7.41(m, 5H), 5.12(s, 2H), 4.36-4.52(m, 3H), 4.02-4.23(m, 6H), 2.63-2.73(m, 1H), 2.56-(dd, J=5, 10Hz, 1H), 1.85-2.12(m, 2H), 1.15-1.23(m, 6H).

EXAMPLE 104N-(2'-Naphthoyl)-(γ-Benzyl ester)-R-Glutamyl-bis(carboethoxymethyl)amide

EDCI (96 mg, 0.50 mmol) was added to a solution of 2-naphthoic acid (87 mg, 0.50 mmol), the product of example 103 (200 mg, 0.45 mmol) and TEA (70 μL, 0.5 mmol) in 5 mL CH₂Cl₂ at room temperature overnight. After evaporation of the solvent, the residue was dissolved in EtOAc and extracted with 0.1 M citric acid(3x) and water(3x). The EtOAc was then dried over MgSO₄, filtered and the solvent evaporated in vacuo. The residue was chromatographed on silica gel eluted with 4:1 to 2:1 step gradient to give 193 mg, 0.34 mmol (76% yield) as an oil. $[\alpha]_D^{25} = +27.7^\circ$ (c=0.73, MeOH). MS(Cl) m/e 563 (m+H)⁺, 374. ¹H NMR (CDCl₃, 300MHz) δ 1.27(dt, J=1.7Hz, 6H), 1.92-2.03(m, 1H), 2.27-2.38(m, 1H), 2.50(dt, J=6.17Hz, 1H), 2.59-2.70-(m, 1H), 4.10(d, J=17Hz, 1H), 4.15-4.23(m, 4H), 4.30(d, J=17Hz, 1H), 4.45(d, J=11Hz, 1H), 4.50(d, J=17Hz, 1H), 5.11(d, J=11Hz, 1H), 5.15(d, J=11Hz, 1H), 5.25-5.32(m, 1H), 7.22(d, J=8Hz, 1H), 7.29-7.35(m, 5H), 7.51-7.60-(m, 2H), 7.82-7.93(m, 4H), 8.32(s, 1H). C,H,N analysis calculated for C₃₁H₃₄N₂O₈ 0.3 H₂O: C 65.55, H 6.14, N 4.93; found C 65.52, H 6.17, N 4.90.

EXAMPLE 105N-(2'-Naphthoyl)-R-Glutamyl-bis(carboethoxymethyl)amide

The product of example 104 (100 mg, 0.18 mmol) was dissolved in 10 mL EtOH and then treated with 25 mg of 10% Pd/C (prewetted with solvent under N₂) under 1 atmosphere H₂ (balloon) overnight. The catalyst was removed by filtration and the solvent was evaporated in vacuo. The residue was redissolved in EtOH and filtered through a 5 μ filter. The filtrate was then mixed with water, frozen and lyophilized to yield an oil 67 mg, (0.14 mmol, 79% yield). R_f = 0.35 (80:20:1 CHCl₃-MeOH-NH₄OH). $[\alpha]_D^{25} = +20.4^\circ$ (c=1.41, MeOH). MS(FAB+) m/e 473(m+H)⁺, 495(m+Na)⁺, 284, 256. ¹H NMR(DMSO-d₆, 300MHz) δ 1.10-(t, J=7Hz, 3H), 1.17(t, J=7Hz, 3H), 1.83-2.08(m, 2H), 2.32(bt, J=8Hz, 2H), 3.86-4.11(m, 5H), 4.19(d, J=17Hz, 1H), 4.48(d, J=18Hz, 1H), 4.55(d, J=18Hz, 1H), 4.43-5.01(m, 1H), 7.57-7.65(m, 2H), 7.93-8.04(m, 4H), 8.52(s, 1H), 8.89(d, J=8Hz, 1H). C,H,N analysis calculated for C₂₄H₂₈N₂O₈ 0.6 H₂O: C 59.65, H 6.09, N 5.80; found C 59.69, H 5.70, N 5.70.

EXAMPLE 106N-(3'-Quinolylcarbonyl)-(γ-Benzyl ester)-R-Glutamyl-bis(carboethoxymethyl)amide

Quinoline-3-carboxylic acid (87 mg, 0.5 mmol) was coupled to the product of example 103 (200 mg, 0.45 mmol) as in example 104. Column chromatography on silica gel eluted with a 9:1 to 2:1 hexane-EtOAc step gradient yielded an oil 131 mg (0.23 mmol, 52% yield). $[\alpha]_D^{25} = +28.0^\circ$ (c=0.55, MeOH). MS(Cl) m/e 564(m+H)⁺. ¹H NMR(CDCl₃, 300MHz) δ 1.23-1.32(m, 6H), 1.92-2.03(m, 1H), 2.28-2.38(m, 1H), 2.51-(dt, J=6.17Hz, 1H), 2.60-2.71(m, 1H), 4.07-4.27(m, 5H), 4.31(d, J=17Hz, 1H), 4.49(s, 2H), 5.13(d, J=11Hz, 1H), 5.18(d, J=11Hz, 1H), 5.25-5.32(m, 1H), 7.28-7.38(m, 6H), 7.62(dt, J=1.8Hz, 1H), 7.82(dt, J=1.8Hz, 1H), 7.90-(dd, J=1.8Hz, 1H), 8.15(d, J=8Hz, 1H), 8.58(d, J=2Hz, 1H), 9.32(d, J=2Hz, 1H). C,H,N analysis calculated for C₃₀H₃₃N₃O₈ 0.5 H₂O: C 62.93, H 5.99, N 7.94; found C 62.95, H 6.09, N 7.22.

EXAMPLE 107

N-(3'-Quinolylcarbonyl)-R-Glutamyl-bis(carboethoxymethyl)amide

The product of example 106 (70 mg, 0.124 mmol) was treated with 20 mg of 10% Pd/C and 1 atmosphere hydrogen gas in 7 mL EtOH. After 6 hours, the catalyst was filtered and the solvent concentrated, mixed with water and then frozen and lyophilized to yield 49 mg (0.10 mmol, 84% yield). mp = 48-52° C. $[\alpha]_D^{25} = +17.1^\circ$ (c=1.37, MeOH). MS(FAB+) m/e 474(m+H)⁺, 496(m+Na)⁺. ¹H NMR-(DMSO-d₆, 300MHz) δ 1.12(t, J=7Hz, 3H), 1.17(t, J=7Hz, 3H), 1.83-2.10(m, 2H), 2.34(bt, J=7Hz, 2H), 3.42-4.12-(m, 5H), 4.19(d, J=17Hz, 1H), 4.51(d, J=17Hz, 1H), 4.55(d, J=17Hz, 1H), 4.95-5.02(m, 1H), 7.71(dt, J=1.8Hz, 1H), 7.88(dt, J=1.8Hz, 1H), 8.10(d, J=8Hz, 2H), 8.88(d, J=2Hz, 1H), 9.18(d, J=7Hz, 1H), 9.28(d, J=2Hz, 1H). C,H,N analysis calculated for C₂₃H₂₇N₃O₈ 0.7 H₂O: C 56.83, H 5.89, N 8.65; found C 56.75, H 5.53, N 8.52.

EXAMPLE 108N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-(γ -Benzyl ester)-R-Glutamyl-bis(carboethoxymethyl)amide

4,8-Dihydroxyquinoline-2-carboxylic acid (103 mg, 0.50 mmol), the product of example 103 (202 mg, 0.45 mmol), HOBt (68 mg, 0.5 mmol) and TEA (70 μ L, 0.5 mmol) were dissolved in 5 mL DMF and treated with EDCI (96 mg, 0.50 mmol) overnight. After evaporation of the solvent, the residue was extracted as in example 82. The residue was chromatographed on silica gel eluted with 1% to 6% EtOH in methylene chloride step gradient to yield 193 mg (0.32 mmol, 72% yield) of nearly pure product. A second silica gel column eluted with 4:1 to 1:3 hexane-EtOAc step gradient yielded pure product. mp = 79-82° C. $[\alpha]_D^{25} = +23.9^\circ$ (c=1.6, MeOH). MS(Cl) m/e 596(m+H)⁺, 407, 379. ¹H NMR(DMSO-d₆, 500MHz) δ 1.18(t, J=6Hz, 3H), 1.23(t, J=6Hz, 3H), 1.98-2.05(m, 1H), 2.13-2.20(m, 1H), 2.37-2.47(m, 2H), 3.82-3.94(m, 2H), 4.00(d, J=16Hz, 1H), 4.04(d, J=7Hz, 1H), 4.07(d, J=7Hz, 1H), 4.18(d, J=16Hz, 1H), 4.38(d, J=17Hz, 1H), 4.53(d, J=17Hz, 1H), 4.98-5.03(m, 1H), 5.05(d, J=11Hz, 1H), 5.08(d, J=11Hz, 1H), 7.08(d, J=8Hz, 1H), 7.28-7.36(m, 4H), 7.42(t, J=7Hz, 1H), 7.52(s, 1H), 7.54(d, J=7Hz, 1H), 7.71(d, J=8Hz, 1H), 10.04(s, 1H), 11.75(bs, 1H). C,H,N analysis calculated for C₃₀H₃₃N₃O₁₀ 0.5 H₂O: C 59.60, H 5.67, N 6.97; found C 59.69, H 5.56, N 6.88.

EXAMPLE 109N-(4',8'-Dihydroxyquinolyl-2-carbonyl)-R-Glutamyl-bis(carboethoxymethyl)amide

The product of example 108 (22 mg, 0.37 mmol) was treated with 5 mg of 10% Pd/C in 5 mL MeOH as in example 97. After 2 hours, the catalyst was filtered and the solvent was evaporated. The crude product was chromatographed on silica gel eluted with 70:30:1 EtOAc-hexanes-HOAc. $[\alpha]_D^{25} = +10.0^\circ$ (c=0.21, MeOH). MS(Cl) m/e 506(m+H)⁺. ¹H NMR(DMSO-d₆, 300MHz) δ 1.07-1.17(m, 6H), 1.86-1.97(m, 1H), 2.04-2.14-(m, 1H), 2.18-2.36(m, 2H), 3.82-3.96(m, 2H), 4.02-4.09(m, 3H), 4.18(d, J=16Hz, 1H), 4.42(d, J=18Hz, 1H), 4.56-(d, J=18Hz, 1H), 4.92-5.00(m, 1H), 7.08(dd, J=1.7Hz, 1H), 7.39(t, J=8Hz, 1H), 7.46(s, 1H), 7.55(dd, J=1.8Hz, 1H), 9.68(d, J=8Hz, 1H), 12.0(bs, 1H). C,H,N analysis calculated for C₂₃H₂₇N₃O₁₀ 1.5 H₂O: C 1.88, H 5.68, N 7.89; found C 52.32, H 5.51, N 7.33.

EXAMPLE 110N-(2'-Indolylcarbonyl)-(N γ -n-pentylamide)-R-Glu-N α ,N α -di-n-pentyl amide

The product of example 4 (50 mg, 0.116 mmol) and n-amyamine (35 μ L, 0.30 mmol) were dissolved in 5 mL CH_2Cl_2 and treated with EDCI (23 mg, 0.12 mmol). TEA (17 μ L, 0.12 mmol) and additional EDCI (23 mg) were added after 1 day. After 2 days, the solvent was evaporated and the residue in EtOAc was washed with 0.1% citric acid, water, dried and evaporated. The crude residue was chromatographed on silica gel eluted with a 4:1 to 1:1 hexane-EtOAc stepped gradient to yield 26.7 mg, 0.054 mmol (46%). mp = 103-5° C. R_f = 0.56(1:1 hexane-EtOAc). MS(Cl) m/e 499(m + H)⁺, 341. ¹H NMR(CDCl_3 , 300MHz) δ 0.92-(t, J = 7Hz, 3H), 1.23-1.38(m, 12H), 1.49-1.60(m, 6H), 1.83-1.95(m, 1H), 2.13-2.23(m, 1H), 2.29-2.33(m, 2H), 3.03-(dt, J = 7, 14Hz, 1H), 3.12-3.47(m, 4H), 3.63(dt, J = 7, 14Hz, 1H), 4.99(dt, J = 2, 9Hz, 1H), 6.73(bt, J = 5Hz, 1H), 7.01-(dd, J = 1, 2Hz, 1H), 7.15(dt, J = 1, 7Hz, 1H), 7.29(dt, J = 1, 7Hz, 1H), 7.42(dd, J = 1, 8Hz, 1H), 7.47(d, J = 8Hz, 1H), 7.66(dd, J = 1, 7Hz, 1H), 9.12(s, 1H). C, H, N analysis calculated for $\text{C}_{29}\text{H}_{46}\text{N}_4\text{O}_3$: C 69.84, H 9.30, N 11.24; found C 69.78, H 9.28, N 11.13.

EXAMPLE 111

N-(2'-Indolylcarbonyl)-[N γ -(2'-phenylethylamide)]-R-Glu-N α , N α -di-n-pentyl amide

The product of example 4 (50 mg, 0.116 mmol) was reacted with β -phenethylamine (38 μ L, 0.30 mmol) in the same manner as in example 110. The reaction yielded 36 mg, 0.068 mmol (58%). MS(Cl) m/e 533-(m + H)⁺, 376, 348. ¹H NMR(CDCl_3 , 300MHz) δ 0.85-0.92(m, 6H), 1.25-1.36(m, 8H), 1.48-1.65(m, 4H), 1.85-1.96(m, 1H), 2.13-2.32(m, 3H), 2.82(t, J = 8Hz, 2H), 3.02-3.22(m, 2H), 3.37-3.65(m, 4H), 5.02(dt, J = 2, 9Hz, 1H), 6.54(t, J = 5Hz, 1H), 6.99(dd, J = 1, 2Hz, 1H), 7.12-7.32(m, 8H), 7.40-7.46(m, 2H), 7.68(dd, J = 1, 7Hz, 1H), 9.11-(s, 1H). C, H, N analysis calculated for $\text{C}_{32}\text{H}_{44}\text{N}_4\text{O}_3 \cdot 0.3 \text{ H}_2\text{O}$: C 71.42, H 8.35, N 10.41; found C 71.46, H 8.32, N 10.43.

EXAMPLE 112

N-(4', 8'-Dihydroxyquinolyl-2'-carbonyl)-(γ -benzyl ester)-R-Glu-di-n-pentylamide

To a solution of the hydrochloride of example 2 (1.0 grams, 2.42 mmol) in DMF (10mL) at 0° C were added NMM (0.55 mL, 5 mmol), HOBT (0.33 gram, 2.42 mmol), 4,8-dihydroxyquinoline-2-carboxylic acid (0.5 gram, 2.44 mmol) and DCC (0.55 grams, 2.6 mmol). The reaction was stirred overnight with warming to ambient temperature. The solvents were evaporated in vacuo and the resulting residue was dissolved in EtOAc. The dicyclohexylurea was filtered away, the filtrate was concentrated and the residue chromatographed on silica gel using CHCl_3 and MeOH (18:1). The solvents were removed and the semisolid yellow residue dried under vacuum at ambient temperature. Product was isolated in 55% yield 0.75 grams. MS(Cl) m/e 564(m + H)⁺. ¹H NMR(CDCl_3 , 300MHz) δ 0.85-0.95(m, 6H), 1.22-1.42(m, 8H), 1.55-1.75(m, 4H), 1.75-2.0-(m, 2H), 2.5-2.62(m, 1H), 2.65-2.8(m, 1H), 3.05-3.20(m, 1H), 3.30-3.52(m, 2H), 3.64-3.78(m, 1H), 5.0(m, 1H), 5.2-(s, 2H), 6.95(m, 2H), 7.15(m, 7H), 7.6(d, J = 9Hz, 1H), 8.1(bs, 1H), 9.2(bs, 1H). C, H, N analysis calculated for $\text{C}_{32}\text{H}_{41}\text{N}_3\text{O}_6 \cdot 0.25 \text{ H}_2\text{O}$: C 67.64, H 7.36, N 7.40; found C 67.62, H 7.03, N 7.66.

EXAMPLE 113

N-(4', 8'-Dihydroxyquinolyl-2'-carbonyl)-R-Glu-N α , N α -di-n-pentyl amide

To a suspension of 0.3 grams 10% Pd/C in MeOH (15 mL) and cyclohexadiene (2 mL) under inert atmosphere (N_2) was added via cannula solution of the benzyl ester of example 112 (0.4 grams, 0.71 mmol)

in MeOH. The reaction mixture was stirred overnight at ambient temperature. The mixture was filtered through celite and washed several times with MeOH. The filtrate and washings were combined and concentrated in vacuo. The product was purified by flash chromatography using EtOAc/hexane and HOAc (70:30:1) as the elutant mixture. Removal of the volatiles in vacuo provided pure product in 60% yield (0.2 grams). MS(Cl) m/e 474(m+H)⁺. ¹H NMR(DMSO-d₆, 300MHz) δ 0.75-0.88(m, 6H), 1.1-1.35(m, 8H), 1.4-1.65-(m, 4H), 1.95-2.10(m, 2H), 2.35-2.42(m, 2H), 3.1(m, 1H), 3.2-3.4(m, 3H), 5.0(m, 1H), 7.10(d, J=9Hz, 1H), 7.42-(t, J=10Hz, 1H), 7.49(bs, 1H), 7.55(d, J=9Hz, 1H), 9.6(bd, J=9Hz, 1H), 10.1(bs, 1H), 12.00(bs, 1H). C,H,N analysis calculated for C₂₅H₃₅N₃O₆ H₂O HOAc: C 58.79, H 7.49, N 7.62; found C 58.78, H 7.04, N 7.87.

EXAMPLE 114

N-(2'-Indolylcarbonyl)-R-Glu-N^γ-methoxyamide-N^α,N^α-di-n-pentylamide

The product of example 4 (50 mg, 0.116 mmol) was reacted with methoxyamine hydrochloride (24 mg, 0.29 mmol) in the same manner as example 110. The reaction yielded 35 mg, 0.076 mmol (66%). ¹H NMR-(CDCl₃, 300MHz) δ 0.89(t, J=7Hz, 6H), 1.23-1.37(m, 8H), 1.48-1.62(m, 4H), 1.85-1.94(m, 1H), 2.20(bs, 1H), 2.24-(bs, 2H), 3.00-3.08(m, 1H), 3.10-3.21(m, 1H), 3.32-3.43(m, 1H), 3.62(dt, J=8, 13Hz, 1H), 3.81(s, 3H), 4.94-(bt, J=8Hz, 1H), 7.02(s, 1H), 7.15(dt, J=1.8Hz, 1H), 7.30(dt, J=1.7Hz, 1H), 7.42(d, J=8Hz, 1H), 7.62-(bd, J=7Hz, 1H), 7.67(d, J=8Hz, 1H), 9.28(s, 1H), 9.94(s, 1H).

EXAMPLE 115

N-Benzyl-(γ-benzyl ester)-R-Glu-N^α,N^α-di-n-pentylamide

The product of example 2 (413 mg, 1.0 mmol) was dissolved in 5 mL of dry ethanol and treated consecutively with TEA (139 μL, 1.0 mmol), benzaldehyde (127 μL, 1.25 mmol) and acetic acid (57 μL, 1.0 mmol) each at 10 minute intervals. Then sodium cyanoborohydride (126 mg, 2.0 mmol) was added in 4 portions over 30 minutes. Bubbling commenced immediately and the reaction became opaque. The reaction was allowed to stir overnight. The reaction was then poured into ethyl acetate, extracted with 0.5 M sodium bicarbonate and then dried over MgSO₄, filtered and the ethyl acetate evaporated in vacuo. The resultant oil was chromatographed on silica gel eluted with a 9:1 to 4:1 step gradient.

EXAMPLE 116

N-Benzyl-N-(3'-quinolylcarbonyl)-R-Glu-N^α,N^α-di-n-pentylamide

The product of example 115, quinoline-3-carboxylic acid and TEA were dissolved in methylene chloride and treated with BOPCI. The residue after evaporation of the solvent was extracted as in example 70 and then purified by column chromatography.

EXAMPLE 117

N-Methyl-N-(3'-quinolylcarbonyl)-(γ-benzyl ester)-R-Glu-N^α,N^α-di-n-pentylamide

The product of example 5 (100 mg, 0.176 mmol) was dissolved in 5 mL of DMF and then treated with lithium bis(trimethylsilyl) amide (170 μ L, 0.17 mmol, 1.0 M in THF) followed by methyl iodide (33 μ L, 0.035 mmol). After 2 hours, the solvents were evaporated and the residue was extracted with 0.1 M citric acid and 0.5 M sodium bicarbonate. Column chromatography on silica gel eluted with 9:1 to 2:1 hexane-ethyl acetate step gradient yielded 48 mg of product (0.089 mmol, 52%).

EXAMPLE 118

N-(3'-Quinolylcarbonyl)-R-Glu-N α ,N α -di-n-pentylamide Dicyclohexylamine salt

The product of example 15 (880 mg, 2 mmol) was dissolved in 10 mL of methylene chloride and then treated with dicyclohexylamine (4 mL, 20 mmol). After complete mixing, hexane was added (200 mL) and a solid formed. The solid was collected and rinsed with fresh hexanes and then dried in vacuo. mp = 129-131 $^{\circ}$ C. MS(Cl) m/e 442(m+H) $^{+}$, 398. 1 H NMR(DMSO-d₆, 300MHz) δ 0.83(d, J=7Hz, 3H), 0.88(d, J=5Hz, 3H), 1.12-1.32(m, 20H), 1.42-1.70(m, 8H), 1.82-1.92(m, 6H), 2.26-2.32(m, 2H), 2.67-2.74(m, 2H), 3.02-(quintet, J=7Hz, 1H), 3.23-3.35(m, 1H), 3.42-3.60(m, 2H), 4.82-4.90(m, 1H), 7.70(dt, J=1.7Hz, 1H), 7.86-(dt, J=1.7Hz, 1H), 8.09(d, J=8Hz, 2H), 8.90(d, J=2Hz, 1H), 9.31(d, J=2Hz, 1H), 9.50(d, J=7Hz, 1H). C, H, N analysis calculated for C₃₇H₅₈N₄O₄: C 71.34, H 9.39, N 9.00; found C 71.50, H 9.40, N 8.91.

The compounds of Formula I antagonize CCK which makes the compounds useful in the treatment and prevention of disease states wherein CCK or gastrin may be involved, for example, gastrointestinal disorders such as irritable bowel syndrome, ulcers, excess pancreatic or gastric secretion, hyperinsulinemia, acute pancreatitis, cancers of the gall bladder and pancreas, motility disorders, pain (potentiation of opiate analgesia), central nervous system disorders caused by CCK's interaction with dopamine such as neuroleptic disorders, tardive dyskinesia, Parkinson's disease, psychosis or Gilles de la Tourette Syndrome, disorders of appetite regulatory systems, Zollinger-Ellison syndrome, and central G cell hyperplasia.

The ability of the compounds of Formula I to interact with CCK receptors and to antagonize CCK can be demonstrated in vitro using the following protocols.

CCK₈ [Asp-Tyr(SO₃H)-Met-Gly-Trp-Met-Asp-Phe-NH₂] was purchased from Beckman Instruments (Palo Alto, CA) and Peptide International (Louisville, KY). Chymostatin, L-Try-Gly, puromycin, bestatin, EGTA, HEPES and BSA were purchased from Sigma Chemical Co. (St. Louis, MO). [¹²⁵I]BH-CCK₈ (specific activity, 2200 Ci/mmol) and Aquasol-2 scintillation cocktail were obtained from New England Nuclear (Boston, MA). Male guinea pigs, 250 to 325 g, were obtained from Scientific Small Animal Laboratory and Farm (Arlington Heights, IL). Collagenase, 300 units per mg, was purchased from Worthington (Frehold, N.J.).

Protocol For Radioligand Binding Experiments

1. Guinea Pig Cerebral Cortical and Pancreatic Membrane Preparations

Cortical and pancreatic membranes were prepared as described (Lin and Miller; J. Pharmacol. Exp. Ther. 232, 775-780, 1985). In brief, cortex and pancreas were removed and rinsed with ice-cold saline. Visible fat and connective tissues were removed from the pancreas. Tissues were weighed and homogenized separately in approximately 25 mL of ice-cold 50 mM Tris-HCl buffer, pH 7.4 at 4 $^{\circ}$ C, with a Brinkman Polytron for 30 sec, setting 7. The homogenates were centrifuged for 10 min at 1075 x g and pellets were discarded. The supernatants were saved and centrifuged at 38,730 x g for 20 min. The resultant pellets were rehomogenized in 25 mL of 50 mM Tris-HCl buffer with a Teflon-glass homogenizer, 5 up and down strokes. The homogenates were centrifuged again at 38,730 x g for 20 min. Pellets were then resuspended in 20 mM HEPES, containing 1 mM EGTA, 118 mM NaCl, 4.7 mM KCl, 5 mM MgCl₂, 100 microM bestatin, 3 microM phosphoramidon, pH 7.4 at 22 $^{\circ}$ C, with a Teflon-glass homogenizer, 15 up and down strokes. Resuspension volume for the cortex was 15-18 mL per gm of original wet weight and 60 mL

per gm for the pancreas.

2. Incubation Conditions

[¹²⁵I]Bolton-Hunter CCK₈, and test compounds were diluted with HEPES-EGTA-salt buffer (see above) containing 0.5% bovine serum albumin (BSA). To 1 mL Skatron polystyrene tubes, were added 25 microliters of test compounds, 25 microliters of [¹²⁵I]BH-CCK₈, and 200 microliters of membrane suspension. the final BSA concentration was 0.1%. The cortical tissues were incubated at 30 °C for 150 min and pancreatic tissues were incubated at 37 °C for 30 min. Incubations were terminated by filtration using Skatron Cell Harvester and SS32 microfiber filter mats. The specific binding of [¹²⁵I]BH-CCK₈, defined as the difference between binding in the absence and presence of 1 microM CCK₈, was 85-90% of total binding in cortex and 90-95% in pancreas. IC₅₀'s were determined from the Hill analysis. The results of these binding assays are shown in Table I.

Protocol for Amylase

This assay was performed using the modified protocol of Lin et al., J. Pharmacol. Exp. Ther, 236, 729-734, 1986.

1. Guinea Pig Acini Preparation

Guinea pig acini were prepared by the method of Bruzzone et al. (Biochem. J. 226, 621-624, 1985) as follows. Pancreas was dissected out and connective tissues and blood vessels were removed. The pancreas was cut into small pieces (2mm) by a scissor and placed in a 15 mL conical plastic tube containing 2.5 mL of Krebs-Ringer HEPES (KRH) buffer plus 400 units per mL of collagenase. The composition of the KRH buffer was: HEPES, 12.5 mM; NaCl, 118 mM; KCl, 4.8 mM; CaCl₂, 1 mM; KH₂PO₄, 1.2 mM; MgSO₄, 1.2 mM; NaHCO₃, 5 mM; glucose, 10 mM, pH 7.4. The buffer was supplemented with 1% MEM vitamins, 1% MEM amino acids and 0.001% aprotinin. The tube was shaken by hand until the suspension appeared homogeneous, usually 5 to 6 min. 5 mL of the KRH, without collagenase and with 0.1% BSA, were added and the tube was centrifuged at 50 x g for 35 sec. The supernatant was discarded and 6 mL of the KRH were added to the cell pellet. Cells were triturated by a glass pipet and centrifuged at 50 x g for 35 sec. This wash procedure was repeated once. The cell pellet from the last centrifugation step was then resuspended in 15 mL of KRH containing 0.1% BSA. The contents were filtered through a dual nylon mesh, size 275 and 75 micrometers. The filtrate, containing the acini, were centrifuged at 50 x g for 3 min. The acini were then resuspended in 5 mL of KRH-BSA buffer for 30 min at 37 °C, under 100% O₂, with a change of fresh buffer at 15 min.

2. Amylase Assay

After the 30 min incubation time, the acini were resuspended in 100 volumes of KRH-BSA buffer, containing 3 microM phosphoramidon and 100 microM bestatin. While stirring, 400 microliters of acini were added to 1.5 mL microcentrifuge tubes containing 50 microliters of CCK₈, buffer, or test compounds. The final assay volume was 500 microliters. Tubes were vortexed and placed in a 37 °C water bath, under 100% O₂, for 30 min. Afterward, tubes were centrifuged at 10,000 g for 1 min. Amylase activity in the supernatant and the cell pellet were separately determined after appropriate dilutions in 0.1% Triton X-100, 10 mM NaH₂PO₄, pH 7.4 by Abbott Amylase A-gent test using the Abbott Bichromatic Analyzer 200. The reference concentration for CCK₈ in determining the IC₅₀'s of the compounds of Formula I was 3x10⁻¹⁰M. The results of this assay are shown in Table 2.

TABLE 1

	Compound of Example	IC ₅₀ (nM)	
		[¹²⁵ I]-BH-CCK ₈ Pancreas	[¹²⁵ I]-BH-CCK ₈ Cortex
5	3	240	3500
	4	20	1400
10	6	62	4200
	8	200	19,000
	10	34	6500
15	12	260	3200
	13	92	5600
	15	5.4	3500
	18	820	2900
20	19	27	2600
	21	210	6600
	22	97	6660
25	23	98	3600
	25	310	>10,000
	29	30	5100
30	34	30	~10,000
	36	200	6300
	37	49	1-10,000
	38	900	7100
35	54	660	>10,000
	60	34	1-10,000
	74	210	2000
40	82	250	10,000
	83	44	>10,000
	84	28	620
	85	100	<10,000
45	89	140	>10,000
	91	68	>10,000
50	93	100	≤10,000
	106	93	~10,000
	107	110	>10,000
55	109	56	620

TABLE 2

Compound of Example	IC ₅₀ (nM) Inhibition of Amylase Release
4	22
6	110
8	590
12	830
15	16
19	55
23	1300
29	130
34	53
74	1200

The results of the assays indicate that the compounds of the invention inhibit specific [¹²⁵I]-BH-CCK-8 receptor binding in the concentration range of 10⁻⁹ to 10⁻⁶ M.

The ability of the compounds of Formula I to interact with CCK receptors and to antagonize CCK can be demonstrated in vivo using the following protocol.

Inhibition Of CCK Induced Gastric Emptying

Three fasted mice were dosed (p.o.) with the test compound. CCK₈ (80 micrograms/kg s.c.) was administered within 60 minutes and charcoal meal (0.1 mL of 10% suspension) was given orally 5 minutes later. The animals were sacrificed within an additional 5 minutes.

Gastric emptying, defined as the presence of charcoal within the intestine beyond the pyloric sphincter, is inhibited by CCK₈. Gastric emptying observed in more than 1 or 3 mice (greater than 1) indicates antagonism of CCK₈.

Gastric emptying was observed with a dose of 100 mg/kg of the compound of example 118 in three out of three mice.

The ability of the compounds of Formula I to antagonize CCK induced hyperinsulinemia can be demonstrated in vivo using the following protocol.

Measurement of Plasma Insulin Level Following Treatment with CCK₈ and a Compound of Formula I

Male mice, 20-30 g, were used in all experiments. The animals were fed with laboratory lab chow and water ad libitum. The compound of Formula I (1 to 100 mg/kg in 0.2 mL of 0.9% saline) was administered i.p. Ten minutes later CCK₈ (0.2 to 200 nmole/kg in 0.2 mL of 0.9% saline) or saline was injected into the tail vein. Two minutes later the animals were sacrificed and blood was collected into 1.5 mL heparinized polypropylene tubes. The tubes were centrifuged at 10,000 x g for 2 min. Insulin levels were determined in the supernatant (plasma) by an RIA method using kits from Radioassay Systems Laboratory (Carson, CA.) or Novo Biolabs (MA.).

The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as loweralkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble

or dispersible products are thereby obtained.

The pharmaceutically acceptable salts of the present invention can be synthesized from the compounds of Formula I which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts are prepared by reacting the free base or acid with stoichiometric amounts or with an excess of the desired salt forming inorganic or organic acid or base in a suitable solvent or various combinations of solvents.

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid. Other salts include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic bases.

The pharmaceutically acceptable salts of the acid of Formula I are also readily prepared by conventional procedures such as treating an acid of Formula I with an appropriate amount of a base, such as an alkali or alkaline earth metal hydroxide e.g. sodium, potassium, lithium, calcium, or magnesium, or an organic base such as an amine, e.g., dibenzylethylenediamine, cyclohexylamine, dicyclohexylamine, trimethylamine, piperidine, pyrrolidine, benzylamine and the like, or a quaternary ammonium hydroxide such as tetramethylammonium hydroxide and the like.

When a compound of Formula I is used as an antagonist of CCK or gastrin in a human subject, the total daily dose administered in single or divided doses may be in amounts, for example, from 0.001 to 1000 mg a day and more usually 1 to 1000 mg. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated, the particular treatment and the particular mode of administration.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease undergoing therapy.

The compounds of the present invention may be administered orally, parenterally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleagenous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.


Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsion, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

The foregoing is merely illustrative of the invention and is not intended to limit the invention to the disclosed compounds. Variations and changes which are obvious to one skilled in the art are intended to be within the scope and nature of the invention which are defined in the appended claims.

Claims

1. A compound of the formula:



wherein

R_1 and R_2 are independently selected from

1) hydrogen,

2) C_1 - C_8 loweralkyl,

3) cycloalkyl,

4) loweralkenyl,

5) $-(CH_2)_m(C(O))_r$ cycloalkyl wherein m and r are not both 0,


6) $-(CH_2)_mCN$,

7) $-(CH_2)_mOR_9$ wherein R_9 is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH (loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,

8) adamantyl,

9) $-(CH_2)_m(C(O))_rNR_6R_7$ wherein R_6 and R_7 are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, and $-(CH_2)_m$ heterocyclic,

10) cyclic groups wherein R_1 and R_2 taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from $(CH_2)_p$, $(CH=CH)_r$, $CH_2C(O)$, $C(O)CH_2$, QCH_2 , CH_2Q , $C(O)Q$, and $QC(O)$ wherein Q is S , O , or NR_{10} wherein R_{10} is hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl), $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$ wherein X is halogen, $-(C(O))_r(CH_2)_sR_{11}$ wherein R_{11} is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ heterocyclic, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as previously defined, $-NR_4OH$ wherein R_4 is hydrogen, $-(C(O))_r$ loweralkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkynyl, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_mOR_4$ wherein R_4 is as defined above, or $-(CH_2)_mSR_4$ wherein R_4 is as defined above; Y is S , O , CH_2 or NR_{10} ; and R_5 is one, two or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, decahydronaphthyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$, $-(C(O))_r(CH_2)_mR_6$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-(CH_2)_mOR_4$, $-(CH_2)_mSR_4$ and $-NR_4OH$ wherein R_4 is as defined above;

- 11) $-(C(R_5)_2)_m$ aryl wherein R_5 is as defined above with the proviso that not all R_5 substituents are hydrogen,
 12) $-(C(R_5)_2)_m$ (substituted aryl) wherein R_5 is defined as above with the proviso that not all R_5 substituents are hydrogen and wherein substituted aryl is as defined above, and
 13) $-(C(R_5)_2)_m C(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

B is

1) $(CH_2)_m$,

2) $(CR_5 = CR_5)_q$ wherein R_5 is as defined above,

3) LCH_2 wherein L is O, S, or NR_5 wherein R_5 is as defined above,

4) CH_2L wherein L is as defined above,

or


5) NR_5 wherein R_5 is as defined above;

D is

1) $C(O)R_3$ wherein R_3 is as defined above

or

2)



Z is

1) $C(O)$,

2) $C(S)$

or

3) $S(O)_2$;

Ar is a heterocyclic group,

n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or

pharmaceutically acceptable salts thereof, with the proviso that when $n=1-3$, D is $-C(O)R_3$ wherein R_3 is

$-OH$ or $-NR_6R_7$ wherein R_6 and R_7 are independently chosen from hydrogen, loweralkyl, cycloalkyl, C_2-4

loweralkenyl, benzyl, phenethyl and naphthylmethyl; R_1 and R_2 are independently chosen from hydrogen, loweralkyl, cycloalkyl, C_2-4 loweralkenyl, benzyl, phenethyl and naphthylmethyl; Z is $C(O)$; and B is a bond

or a C_1-5 alkylene, C_2-5 alkenylene or C_4-6 alkadienylene group; then Ar is not thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, im-

idazoliny, pyrazolidinyl, triazolyl, tetrazolyl, tetrahydrofuryl, pyranyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, piperidinyl, piperazinyl, morpholinyl, triazinyl, thiazinyl, benzothienyl, benzofuryl, isobenzofuryl,

indolyl, isoindolyl, indazolyl, purinyl, quinolyl, isoquinolyl, naphthyridinyl, quinoxalinyl, chromanyl, indolinyl, isoindolinyl, or chromenyl, unsubstituted or substituted by at least one substituent selected from halogen,

hydroxyl, amino, carboxyl, C_1-8 alkyl, oxo, C_1-8 alkoxy, phenyl, naphthyl, chlorophenyl, benzyl, phenethyl, naphthylmethyl, (2,3,4- or 3,4,5-)trimethoxybenzyl, amino- C_1-4 alkyl, di- C_1-4 alkylamino- C_1-4 alkyl, hydroxy-

substituted C_1-4 alkyl, C_1-5 alkanoyloxy, styrylcarbonyl, 3,4,5-trimethoxystyrylcarbonyl, benzyloxycarbonylamino, benzyloxycarbonylaminomethyl, benzoylamino, 3,4,5-trimethoxybenzoylamino, pyrrolidinylcar-


bonylmethyl, piperidylcarbonylmethyl, piperazinylcarbonylmethyl and morpholinylcarbonylmethyl.

2. The compound of Claim 1 wherein D is $-C(O)R_3$ wherein R_3 is defined above, Z is $C(O)$, and Ar is

indolyl or quinolyl, or pharmaceutically acceptable salts thereof.

3. A compound selected from the group consisting of N-(2-indolylcarbonyl)-R-Glu-(di-n-pentyl) amide, N-(3'-quinolylcarbonyl)-R-Glu-(di-n-pentyl) amide and Ethyl N-(3'-quinolylcarbonyl)-R-Glutamyl(N-benzyl)-glyconate, or pharmaceutically acceptable salts thereof.


4. A pharmaceutical composition useful for antagonizing CCK, comprising a pharmaceutical carrier and a therapeutically effective amount of a compound of the formula:



wherein

R₁ and R₂ are independently selected from

- 1) hydrogen,
- 2) C₁-C₈ loweralkyl,
- 3) cycloalkyl,
- 4) loweralkenyl,
- 5) -(CH₂)_m(C(O))_rcycloalkyl wherein m and r are not both 0,
- 6) -(CH₂)_mCN,
- 7) -(CH₂)_mOR₉ wherein R₉ is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH(loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,
- 8) adamantyl,
- 9) -(CH₂)_m(C(O))_rNR₆R₇ wherein R₆ and R₇ are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, -(CH₂)_maryl, -(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, and --(CH₂)_mheterocyclic,
- 10) cyclic groups wherein R₁ and R₂ taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from (CH₂)_p, (CH=CH)_r, CH₂C(O), C(O)CH₂, QCH₂, CH₂Q, C(O)Q, and QC(O) wherein Q is S, O, or NR₁₀ wherein R₁₀ is hydrogen, -(C(O))_r(C₁-C₈loweralkyl), -(C(O))_rcycloalkyl, -(C(O))_rloweralkenyl, -(C(O))_rloweralkynyl, thioalkoxy, -(C(O))_r(CH₂)_maryl, -(C(O))_r(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mcarboxyl, -(CH₂)_s(C(O))_raryl, -(CH₂)_s(C(O))_r(substituted aryl), -(CH₂)_mcarboalkoxy, -(CH₂)_mcarboxamide, -(CH₂)_mcarboaryloxy, -C(X)₃ wherein X is halogen, -(C(O))_r(CH₂)_sR₁₁ wherein R₁₁ is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, -(CH₂)_maryl, --(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mheterocyclic, nitro, halogen, -CH, -OH, -NH₂, -NHOH, -NR₆R₇ wherein R₆ and R₇ are as previously defined, -NR₄OH wherein R₄ is hydrogen, -(C(O))_rloweralkyl, -(C(O))_rloweralkenyl, -(C(O))_rcycloalkyl, -(C(O))_rloweralkynyl, -(C(O))_r(CH₂)_maryl, -(C(O))_r(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mOR₄ wherein R₄ is as defined above, or -(CH₂)_mSR₄ wherein R₄ is as defined above; Y is S, O, CH₂ or NR₁₀; and R₅ is one, two or three substituents independently selected from hydrogen, -(C(O))_r(C₁-C₈loweralkyl), -(C(O))_rcycloalkyl, decahydronaphthyl, -(C(O))_rloweralkenyl, -(C(O))_rloweralkynyl, thioalkoxy, -(C(O))_r(CH₂)_maryl, -(C(O))_r(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_s(C(O))_raryl, -(CH₂)_s(C(O))_r(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mcarboxyl, -(CH₂)_mcarboalkoxy, -(CH₂)_mcarboxamide, -(CH₂)_mcarboaryloxy, -C(X)₃, -(C(O))_r(CH₂)_mR₆, nitro, halogen, -CN, -OH, -NH₂, -NHOH, -NR₆R₇ wherein R₆ and R₇ are as defined above, -(CH₂)_mOR₄, -(CH₂)_mSR₄ and -NR₄OH wherein R₄ is as defined above;

11) -(C(R₅)₂)_maryl wherein R₅ is as defined above with the proviso that not all R₅ substituents are hydrogen,

12) -(C(R₅)₂)_m(substituted aryl) wherein R₅ is defined as above with the proviso that not all R₅ substituents

are hydrogen and wherein substituted aryl is as defined above, and

13) $-(C(R_5)_2)_mC(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

5 B is

1) $(CH_2)_m$,

2) $(CR_5 = CR_5)_q$ wherein R_5 is as defined above,

3) LCH_2 wherein L is O , S , or NR_5 wherein R_5 is as defined above,

4) CH_2L wherein L is as defined above,

10 or


5) NR_5 wherein R_5 is as defined above;

D is

1) $C(O)R_3$ wherein R_3 is as defined above

or

15 2)



20

Z is

25 1) $C(O)$,

2) $C(S)$


or

3) $S(O)_2$;

Ar is a heterocyclic group,

30 n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or pharmaceutically acceptable salts thereof.

5. A method for antagonizing CCK comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of the formula:



35

40

45

wherein

R_1 and R_2 are independently selected from

1) hydrogen,

2) C_1-C_8 loweralkyl,

50 3) cycloalkyl,

4) loweralkenyl,

5) $-(CH_2)_mC(O)$, cycloalkyl wherein m and r are not both 0,

6) $-(CH_2)_mCN$,


7) $-(CH_2)_mOR_9$ wherein R_9 is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH (loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,

55

8) adamantyl,

9) $-(CH_2)_m(C(O))_rNR_6R_7$ wherein R_6 and R_7 are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, and $-(CH_2)_m$ heterocyclic,

10) cyclic groups wherein R_1 and R_2 taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from $(CH_2)_p$, $(CH=CH)_r$, $CH_2C(O)$, $C(O)CH_2$, QCH_2 , CH_2Q , $C(O)Q$, and $QC(O)$ wherein Q is S , O , or NR_{10} wherein R_{10} is hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl), $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$ wherein X is halogen, $-(C(O))_r(CH_2)_sR_{11}$ wherein R_{11} is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ heterocyclic, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as previously defined, $-NR_4OH$ wherein R_4 is hydrogen, $-(C(O))_r$ loweralkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkynyl, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_mOR_4$ wherein R_4 is as defined above, or $-(CH_2)_mSR_4$ wherein R_4 is as defined above; Y is S , O , CH_2 or NR_{10} ; and R_5 is one, two or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, decahydronaphthyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$, $-(C(O))_r(CH_2)_mR_6$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-(CH_2)_mOR_4$, $-(CH_2)_mSR_4$ and $-NR_4OH$ wherein R_4 is as defined above;

11) $-(C(R_5)_2)_m$ aryl wherein R_5 is as defined above with the proviso that not all R_5 substituents are hydrogen, 12) $-(C(R_5)_2)_m$ (substituted aryl) wherein R_5 is defined as above with the proviso that not all R_5 substituents are hydrogen and wherein substituted aryl is as defined above, and

13) $-(C(R_5)_2)_mC(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

B is

1) $(CH_2)_m$,

2) $(CR_5=CR_5)_q$ wherein R_5 is as defined above,

3) LCH_2 wherein L is O , S , or NR_5 wherein R_5 is as defined above,

4) CH_2L wherein L is as defined above,

or


5) NR_5 wherein R_5 is as defined above;

D is

1) $(C(O)R_3)$ wherein R_3 is as defined above

or

2)



Z is

1) $C(O)$,

2) C(S)


or

3) S(O)₂;

Ar is a heterocyclic group,

5 n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or pharmaceutically acceptable salts thereof.

6. A pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount useful for treatment or prevention of disorders of the gastrointestinal, central nervous, appetite regulating or pain regulating systems of a compound of the formula:



wherein

R₁ and R₂ are independently selected from

1) hydrogen,

2) C₁-C₈ loweralkyl,

3) cycloalkyl,

4) loweralkenyl,


5) -(CH₂)_m(C(O))_rcycloalkyl wherein m and r are not both 0,6) -(CH₂)_mCN,

7) -(CH₂)_mOR₉ wherein R₉ is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino and NH(loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,

8) adamantyl,

9) -(CH₂)_m(C(O))_rNR₆R₇ wherein R₆ and R₇ are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, -(CH₂)_maryl, -(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above; and --(CH₂)_mheterocyclic,

10) cyclic groups wherein R₁ and R₂ taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from (CH₂)_p, (CH=CH)_r, CH₂C(O), C(O)CH₂, QCH₂, CH₂Q, C(O)Q, and QC(O) wherein Q is S, O, or NR₁₀ wherein R₁₀ is hydrogen, -(C(O))_r(C₁-C₈loweralkyl), -(C(O))_rcycloalkyl, -(C(O))_rloweralkenyl, -(C(O))_rloweralkynyl, thioalkoxy, -(C(O))_r(CH₂)_maryl, -(C(O))_r(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mcarboxyl, -(CH₂)_s(C(O))_raryl, -(CH₂)_s(C(O))_r(substituted aryl), -(CH₂)_mcarboalkoxy, -(CH₂)_mcarboxamide, -(CH₂)_mcarboaryloxy, -C(X)₃ wherein X is halogen, -(C(O))_r(CH₂)_sR₁₁ wherein R₁₁ is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, -(CH₂)_maryl, --(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mheterocyclic, nitro, halogen, -CN, -OH, -NH₂, -NHOH, -NR₆R₇ wherein R₆ and R₇ are as previously defined, -NR₄OH wherein R₄ is hydrogen, -(C(O))_rloweralkyl, -(C(O))_rloweralkenyl, -(C(O))_rcycloalkyl, -(C(O))_rloweralkynyl, -(C(O))_r(CH₂)_maryl, -(C(O))_r(CH₂)_m(substituted aryl) wherein substituted aryl is as defined above, -(CH₂)_mOR₄ wherein R₄ is as defined above, or -(CH₂)_mSR₄ wherein R₄ is as defined above; Y is S, O, CH₂ or NR₁₀; and R₅ is one, two

or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_8\text{loweralkyl})$, $-(C(O))_r\text{cycloalkyl}$, decahydronaphthyl, $-(C(O))_r\text{loweralkenyl}$, $-(C(O))_r\text{loweralkynyl}$, thioalkoxy, $-(C(O))_r(CH_2)_m\text{aryl}$, $-(C(O))_r(CH_2)_m\text{-(substituted aryl)}$ wherein substituted aryl is as defined above, $-(CH_2)_s(C(O))\text{aryl}$, $-(CH_2)_s(C(O))\text{-(substituted aryl)}$ wherein substituted aryl is as defined above, $-(CH_2)_m\text{carboxyl}$, $-(CH_2)_m\text{carboalkoxy}$, $-(CH_2)_m\text{carboxamide}$, $-(CH_2)_m\text{carboaryloxy}$, $-C(X)_3$, $-(C(O))_r(CH_2)_mR_6$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-(CH_2)_mOR_4$, $-(CH_2)_mSR_4$ and $-NR_4OH$ wherein R_4 is as defined above;

11) $-(C(R_5)_2)_m\text{aryl}$ wherein R_5 is as defined above with the proviso that not all R_5 substituents are hydrogen,

12) $-(C(R_5)_2)_m\text{-(substituted aryl)}$ wherein R_5 is defined as above with the proviso that not all R_5 substituents are hydrogen and wherein substituted aryl is as defined above, and

13) $-(C(R_5)_2)_mC(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

B is

1) $(CH_2)_m$,

2) $(CR_5=CR_5)_q$ wherein R_5 is as defined above,

3) LCH_2 wherein L is O , S , or NR_5 wherein R_5 is as defined above,

4) CH_2L wherein L is as defined above,

or


5) NR_5 wherein R_5 is as defined above;

D is

1) $C(O)R_3$ wherein R_3 is as defined above

or

2)



Z is

1) $C(O)$,

2) $C(S)$

or

3) $S(O)_2$;

Ar is a heterocyclic group,

n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or pharmaceutically acceptable salts thereof,

with the proviso that when the disorder being treated or prevented is ulcers and when $n=1-3$, D is $-C(O)R_3$

wherein R_3 is $-OH$ or $-NR_6R_7$ wherein R_6 and R_7 are independently chosen from hydrogen, loweralkyl,

cycloalkyl, C_{2-4} loweralkenyl, benzyl, phenethyl and naphthylmethyl; R_1 and R_2 are independently chosen

from hydrogen, loweralkyl, cycloalkyl, C_{2-4} loweralkenyl, benzyl, phenethyl and naphthylmethyl; Z is $C(O)$;

and B is a bond or a C_{1-5} alkylene, C_{2-5} alkenylene or C_{4-6} alkadienylene group; then Ar is not thienyl,

furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolidinyl, pyrrolinyl, im-

idazolidinyl, imidazolinyl, pyrazolidinyl, triazolyl, tetrazolyl, tetrahydrofuryl, pyranyl, pyridyl, pyrazinyl,

pyrimidinyl, pyridazinyl, piperidyl, piperazinyl, morpholinyl, triazinyl, thiazinyl, benzothienyl, benzofuryl,

isobenzofuryl, indolyl, isoindolyl, indazolyl, purinyl, quinolyl, isoquinolyl, naphthyridinyl, quinoxadinyl,

chromanyl, indolinyl, isoindolinyl, or chromenyl, unsubstituted or substituted by at least one substituent

selected from halogen, hydroxyl, amino, carboxyl, C_{1-8} alkyl, oxo, C_{1-8} alkoxy, phenyl, naphthyl,

chlorophenyl, benzyl, phenethyl, naphthylmethyl, (2,3,4- or 3,4,5-trimethoxybenzyl, amino- C_{1-4} alkyl, di-


C_{1-4} alkylamino- C_{1-4} alkyl, hydroxy-substituted C_{1-4} alkyl, C_{1-5} alkanoyloxy, styrylcarbonyl, 3,4,5-trimethox-

ystyrylcarbonyl, benzyloxycarbonylamino, benzyloxycarbonylaminomethyl, benzoylamino, 3,4,5-trimethox-

benzoylamino, pyrrolidinylcarbonylmethyl, piperidylcarbonylmethyl, piperazinylcarbonylmethyl and mor-

pholinylcarbonylmethyl.


7. A method for treatment of prevention of disorders of the gastrointestinal, central nervous, appetite regulating a pain regulating systems comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of the formula:



wherein

R_1 and R_2 are independently selected from

- 1) hydrogen,
- 2) C_1 - C_8 loweralkyl,
- 3) cycloalkyl,
- 4) loweralkenyl,
- 5) $-(CH_2)_m(C(O))_r$ cycloalkyl wherein m and r are not both 0,
- 6) $-(CH_2)_mCN$,
- 7) $-(CH_2)_mOR_3$ wherein R_3 is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH (loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,
- 8) adamantyl,
- 9) $-(CH_2)_m(C(O))_rNR_6R_7$ wherein R_6 and R_7 are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, and $-(CH_2)_m$ heterocyclic,
- 10) cyclic groups wherein R_1 and R_2 taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from $(CH_2)_p$, $(CH=CH)_n$, $CH_2C(O)$, $C(O)CH_2$, $QCCH_2$, CH_2Q , $C(O)Q$, and $QC(O)$ wherein Q is S , O , or NR_{10} wherein R_{10} is hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl), $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$ wherein X is halogen, $-(C(O))_r(CH_2)_sR_{11}$ wherein R_{11} is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ heterocyclic, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as previously defined, $-NR_4OH$ wherein R_4 is hydrogen, $-(C(O))_r$ loweralkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkynyl, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_mOR_4$ wherein R_4 is as defined above, $-(CH_2)_mSR_4$ wherein R_4 is as defined above; Y is S , O , CH_2 or NR_{10} ; and R_5 is one, two or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, decahydronaphthyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$, $-(C(O))_r(CH_2)_mR_5$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-(CH_2)_mOR_4$, $-(CH_2)_mSR_4$ and $-NR_4OH$ wherein R_4 is as defined above;

- 11) $-(C(R_5)_2)_m$ aryl wherein R_5 is as defined above with the proviso that not all R_5 substituents are hydrogen,
- 12) $-(C(R_5)_2)_m$ (substituted aryl) wherein R_5 is defined as above with the proviso that not all R_5 substituents

are hydrogen and wherein substituted aryl is as defined above, and

13) $-(C(R_5)_2)_mC(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

5 B is

1) $(CH_2)_m$,

2) $(CR_5 = CR_5)_q$ wherein R_5 is as defined above,

3) LCH_2 wherein L is O, S, or NR_5 wherein R_5 is as defined above,

4) CH_2L wherein L is as defined above,

10 or


5) NR_5 wherein R_5 is as defined above;

D is

1) $C(O)R_3$ wherein R_3 is as defined above

or

15 2)



20

Z is

25 1) $C(O)$,

2) $C(S)$

or

3) $S(O)_2$;


Ar is a heterocyclic group,

30 n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or pharmaceutically acceptable salts thereof, with the proviso that when the disorder being treated or prevented is ulcers and when $n = 1-3$, D is $-C(O)R_3$ wherein R_3 is $-OH$ or $-NR_6R_7$ wherein R_6 and R_7 are independently chosen from hydrogen, loweralkyl, cycloalkyl, C_2-4 loweralkenyl, benzyl, phenethyl and naphthylmethyl; R_1 and R_2 are independently chosen from hydrogen, loweralkyl, cycloalkyl, C_2-4 -loweralkenyl, benzyl, phenethyl and naphthylmethyl; Z is $C(O)$;
35 and B is a bond or a C_1-5 alkylene, C_2-5 -alkenylene or C_4-6 alkadienylene group; then Ar is not thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolynyl, pyrazolidinyl, triazolyl, tetrazolyl, tetrahydrofuryl, pyranyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, piperidyl, piperazinyl, morpholinyl, triazinyl, thiazinyl, benzothienyl, benzofuryl, isobenzofuryl, indolyl, isoindolyl, indazolyl, purinyl, quinolyl, isoquinolyl, naphthyridinyl, quinoxadinyl,
40 chromanyl, indolinyl, isoindolinyl, or chromenyl, unsubstituted or substituted by at least one substituent selected from halogen, hydroxyl, amino, carboxyl, C_1-8 alkyl, oxo, C_1-8 alkoxy, phenyl, naphthyl, chlorophenyl, benzyl, phenethyl, naphthylmethyl, (2,3,4- or 3,4,5-)trimethoxybenzyl, amino- C_1-4 alkyl, di- C_1-4 alkylamino- C_1-4 alkyl, hydroxy-substituted C_1-4 alkyl, C_1-5 alkanoyloxy, sterylcarbonyl, 3,4,5-trimethoxy-
45 styrylcarbonyl, benzyloxycarbonylamino, benzyloxycarbonylaminomethyl, benzoylamino, 3,4,5-trimethoxybenzoylamino, pyrrolidinylcarbonylmethyl, piperidylcarbonylmethyl, piperazinylcarbonylmethyl and morpholinylcarbonylmethyl.

8. A pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount useful for treatment or prevention of hyperinsulinemia of a compound of the formula:

50


55



wherein

R_1 and R_2 are independently selected from

- 1) hydrogen,
- 2) C_1 - C_8 loweralkyl,
- 3) cycloalkyl,
- 4) loweralkenyl,
- 5) $-(CH_2)_m(C(O))_r$ cycloalkyl wherein m and r are not both 0,
- 6) $-(CH_2)_mCN$,
- 7) $-(CH_2)_mOR_9$ wherein R_9 is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH (loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,
- 8) adamantyl,
- 9) $-(CH_2)_m(C(O))_rNR_6R_7$ wherein R_6 and R_7 are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, and $-(CH_2)_m$ heterocyclic,
- 10) cyclic groups wherein R_1 and R_2 taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from $(CH_2)_p$, $(CH=CH)_r$, $CH_2C(O)$, $C(O)CH_2$, QCH_2 , CH_2Q , $C(O)Q$, and $QC(O)$ wherein Q is S , O , or NR_{10} wherein R_{10} is hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_rcycloalkyl$, $-(C(O))_rloweralkenyl$, $-(C(O))_rloweralkynyl$, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl), $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$ wherein X is halogen, $-(C(O))_r(CH_2)_sR_{11}$ wherein R_{11} is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ heterocyclic, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as previously defined, $-NR_4OH$ wherein R_4 is hydrogen, $-(C(O))_rloweralkyl$, $-(C(O))_rloweralkenyl$, $-(C(O))_rcycloalkyl$, $-(C(O))_rloweralkynyl$, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_mOR_4$ wherein R_4 is as defined above, or $-(CH_2)_mSR_4$ wherein R_4 is as defined above; Y is S , O , CH_2 or NR_{10} ; and R_5 is one, two or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_rcycloalkyl$, decahydronaphthyl, $-(C(O))_rloweralkenyl$, $-(C(O))_rloweralkynyl$, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_s(C(O))_r$ aryl, $-(CH_2)_s(C(O))_r$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$, $-(C(O))_r(CH_2)_mR_6$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-(CH_2)_mOR_4$, $-(CH_2)_mSR_4$ and $-NR_4OH$ wherein R_4 is as defined above;

- 11) $-(C(R_5)_2)_m$ aryl wherein R_5 is as defined above with the proviso that not all R_5 substituents are hydrogen,
- 12) $-(C(R_5)_2)_m$ (substituted aryl) wherein R_5 is defined as above with the proviso that no all R_5 substituents

are hydrogen and wherein substituted aryl is as defined above, and

13) $-(C(R_5)_2)_mC(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

B is

1) $(CH_2)_m$,

2) $(CR_5=CR_5)_q$ wherein R_5 is as defined above,

3) LCH_2 wherein L is O , S , or NR_5 wherein R_5 is as defined above,

4) CH_2L wherein L is as defined above,

or


5) NR_5 wherein R_5 is as defined above;

D is

1) $C(O)R_3$ wherein R_3 is as defined above

or

2)



Z is

1) $C(O)$,

2) $C(S)$


or

3) $S(O)_2$;

Ar is a heterocyclic group,

n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or pharmaceutically acceptable salts thereof.

9. A method for treatment or prevention of hyperinsulinemia comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of the formula:



wherein

R_1 and R_2 are independently selected from

1) hydrogen,

2) C_1 - C_8 loweralkyl,

3) cycloalkyl,

4) loweralkenyl,

5) $-(CH_2)_m(C(O))_r$ cycloalkyl wherein m and r are not both 0,


6) $-(CH_2)_mCN$,

7) $-(CH_2)_mOR_3$ wherein R_3 is loweralkyl, aryl, substituted aryl wherein the aryl group is substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, thioalkoxy, carboxy, carboalkoxy, nitro, trihalomethyl, hydroxy, amino, and NH (loweralkyl), arylalkyl or (substituted aryl)alkyl wherein substituted aryl is as defined above,

8) adamantyl,

9) $-(CH_2)_m(C(O))_rNR_6R_7$ wherein R_6 and R_7 are independently selected from hydrogen, loweralkyl, cycloalkyl, loweralkenyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, and $-(CH_2)_m$ heterocyclic,

10) cyclic groups wherein R_1 and R_2 taken together with the adjacent nitrogen atom are morpholinyl, pyrrolidinyl, piperazinyl, piperidinyl or other substituted cyclic groups represented by



wherein E and J are independently selected from $(CH_2)_p$, $(CH=CH)_r$, $CH_2C(O)$, $C(O)CH_2$, QCH_2 , CH_2Q , $C(O)Q$, and $QC(O)$ wherein Q is S, O, or NR_{10} wherein R_{10} is hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_5(C(O))_r$ aryl, $-(CH_2)_5(C(O))_r$ (substituted aryl), $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$ wherein X is halogen, $-(C(O))_r(CH_2)_5R_{11}$ wherein R_{11} is hydrogen, loweralkyl, loweralkenyl, cycloalkyl, $-(CH_2)_m$ aryl, $-(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ heterocyclic, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as previously defined, $-NR_4OH$ wherein R_4 is hydrogen, $-(C(O))_r$ loweralkyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ cycloalkyl, $-(C(O))_r$ loweralkynyl, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_mOR_4$ wherein R_4 is as defined above, or $-(CH_2)_mSR_4$ wherein R_4 is as defined above; Y is S, O, CH_2 or NR_{10} ; and R_5 is one, two or three substituents independently selected from hydrogen, $-(C(O))_r(C_1-C_8$ loweralkyl), $-(C(O))_r$ cycloalkyl, decahydronaphthyl, $-(C(O))_r$ loweralkenyl, $-(C(O))_r$ loweralkynyl, thioalkoxy, $-(C(O))_r(CH_2)_m$ aryl, $-(C(O))_r(CH_2)_m$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_5(C(O))_r$ aryl, $-(CH_2)_5(C(O))_r$ (substituted aryl) wherein substituted aryl is as defined above, $-(CH_2)_m$ carboxyl, $-(CH_2)_m$ carboalkoxy, $-(CH_2)_m$ carboxamide, $-(CH_2)_m$ carboaryloxy, $-C(X)_3$, $-(C(O))_r(CH_2)_mR_6$, nitro, halogen, $-CN$, $-OH$, $-NH_2$, $-NHOH$, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-(CH_2)_mOR_4$, $-(CH_2)_mSR_4$ and $-NR_4OH$ wherein R_4 is as defined above;

11) $-(C(R_5)_2)_m$ aryl wherein R_5 is as defined above with the proviso that not all R_5 substituents are hydrogen,

12) $-(C(R_5)_2)_m$ (substituted aryl) wherein R_5 is defined as above with the proviso that not all R_5 substituents are hydrogen and wherein substituted aryl is as defined above, and


13) $-(C(R_5)_2)_mC(O)R_3$, wherein R_5 is as defined above and wherein R_3 is $-OH$, $-OR_4$ wherein R_4 is as defined above, $-NR_6R_7$ wherein R_6 and R_7 are as defined above, $-NHR_4$ wherein R_4 is as defined above, or $-NR_4OH$ wherein R_4 is as defined above;

B is

- 1) $(CH_2)_m$,
- 2) $(CR_5 = CR_5)_q$ wherein R_5 is as defined above,
- 3) LCH_2 wherein L is O, S, or NR_5 wherein R_5 is as defined above,
- 4) CH_2L wherein L is as defined above,
- or
- 5) NR_5 wherein R_5 is as defined above;

D is

- 1) $C(O)R_3$ wherein R_3 is as defined above
- or
- 2)



Z is

- 1) $C(O)$,

2) C(S)

or

3) S(O)₂;

Ar is a heterocyclic group,

5 n is 1 to 3, m is 0 to 4, p is 0 to 2, r is 0 to 1 and s is 1 to 4; or pharmaceutically acceptable salts thereof.

10

15

20

25

30

35

40

45

50

55



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 88115462.9
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
D,A	US - A - 4 610 983 (NOBORU TAKA-GAWA et al.) * Abstract; columns 15,16,19, 20,31-52 * --	1,6	C 07 D 209/42 C 07 D 213/56 C 07 D 215/16
D,A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 78, no. 10, October 1981 W.F.HAHNE et al. "Proglumide and benzotript: Members of a different class of cholecystokinin receptor antagonists" pages 6304-6308 * Page 6304; abstract * --	1,4,6	C 07 D 241/44 C 07 D 403/12 A 61 K 31/395
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 D 209/00 C 07 D 213/00 C 07 D 215/00 C 07 D 241/00 C 07 D 403/00
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-4,6,8 Claims searched incompletely: - Claims not searched: 5,7,9 Reason for the limitation of the search:</p> <p>(Method for treatment of the human or animal body by therapy (Art. 52(4) EPC))</p>			
Place of search VIENNA		Date of completion of the search 21-12-1988	Examiner HEIN
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO Form 1505.1: 03.82



EP 88115462.9

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D,A	<p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 83, no. 13, July 1986</p> <p>BEN E.EVANS et al. "Design of potent, orally effective, nonpeptidal antagonists of the peptide hormone cholecystokinin"</p> <p>pages 4918-4922</p> <p>* Page 4918; abstract; page 4920, table 2 *</p> <p>-----</p>	1,4,6	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)